

Comparison of Clinical Function Outcomes of Patients Performed with Cemented and Uncemented Bipolar Hemiarthroplasty Surgery on Femoral Neck Fractures Using Modified Harris Hip Score and Proportion of Dislocations after Hip Joint Replacement Surgery in Medan

Stefan Andika^a, Pranajaya Dharma Kadar^a, Muhammad Hidayat Siregar^a

^aDepartment of Orthopaedic and Traumatology, Faculty of Medicine Universitas Sumatera Utara, Adam Malik Hospital, Medan, Indonesia

Abstract

Introduction

Fractures of the neck femur that occur in the elderly are caused by minor trauma with most common age more than 50 years old. While in young adult patients, fractures of the neck femur are caused by severe trauma such as a fall from a height or a traffic accident. Hemiarthroplasty cemented or uncemented are the most common methods of treatment neck femur fracture in elderly patients. Modified Harris Hip Score (MHHS) is a tool to measure the functional capacity and pain to assess the conditions of the patients before and after surgical procedures

Methodology

This is an observational analytic research study with a case control design at Adam Malik Hospital Medan and several network hospitals in Medan. Total of 32 patients with age more than 50 years old were diagnosed with femoral neck fracture with 16 patients underwent cemented bipolar hemiarthroplasty and 16 patients underwent uncemented bipolar hemiarthroplasty surgery during the period September 2021 until August 2022 assessed clinical functional outcome with MHHS scores and the proportion of dislocation 1 month postoperative.

Result

Mean MHHS value for cemented bipolar hemiarthroplasty was 79.00 ± 5.40 and uncemented bipolar hemiarthroplasty 78.81 ± 7.43 and there were no dislocations 1 month post operation.

Discussion

Statistical analytic using independent T-Test showing there was no significant different between MHHS scores in cemented bipolar arthroplasty and uncemented bipolar hemiarthroplasty 1 months post operative with P Value 0.935 (>0.05) without any dislocations was found.

Keywords: Neck Femur Fracture, Modified Harris Hip Score, Bipolar Hemiarthroplasty

1. Introduction

Hip fracture is the most common injury and is found in geriatric patients so it can affect the health system as a whole. Despite the lack of support from the development of implants, surgical techniques and patient care, hip fractures still have a major impact on the existing health system. As many as 250,000 hip fractures in America occur each year and this number will double by 2050 as the geriatric population increases. With the development of the world of science and technology, the mobility of the Indonesian people is also getting higher and this can be seen by the increasing number of motorized vehicle users in Indonesia. (Bucholz et al, 2020).

The hip fracture mortality rate increases by 1 person in 1 year with a mortality range of around 14-36%. According to the journal that has been studied, one of the problems with fractures in the proximal femur is the patient's ability to return to activities before the trauma. 50% of these patients require assistance in carrying out their daily activities and as many as 25% require assistance in the long term to carry out daily activities. In addition, the type of proximal femoral fracture injury also has a significant increase in areas that have a trauma center. (Bucholz et al, 2020; Blomfeldt et al, 2007, p160-165)

Femoral neck and intertrochanteric fractures of the femur are the most common fractures in the elderly. This fracture is closely related to osteoporosis that occurs in the elderly. Fractures of the hip bone also often

occur at a young age that are not associated with osteoporosis but are associated with traffic accidents or falls from a height. (Dilogo IH et al, 2013, p1-15).

Arthroplasty has then become the method of choice in the majority of fractures of the hip region and in several prospective studies it has shown a better outcome and a lower ratio for reoperation. In old age hemiarthroplasty is more considered because it provides a shorter operating time, a simpler surgical technique and a lower risk of dislocation.

However, there is a risk of loss of the acetabular cartilage surface due to friction with metal material from the prosthetic femoral head so that in the future it requires conversion to a total hip arthroplasty. Several risk factors for acetabular joint cartilage erosion are young age, high activity and long follow-up period. (Blomfeldt et al, 2007, p160-165).

In a study conducted by Gupta et al in 2019 in India, cemented bipolar hemiarthroplasty was performed on 9 patients and uncemented bipolar hemiarthroplasty in 10 patients with a diagnosis of femoral neck fracture, evaluated for 6 weeks, 12 weeks and 1 year. At 6 weeks postoperatively there was no significant difference, but at 1 year after surgery, the clinical outcome of MHHS in the cemented group was better than the uncemented group. (Gupta et al, 2019, p622-626).

2. Experimental Methodology

This research is an observational analytic with case control design, to evaluate Comparison of Clinical Function Outcomes of Patients Performed with Cemented and Uncemented Bipolar Hemiarthroplasty Surgery on Femoral Neck Fractures Using Modified Harris Hip Score and Proportion of Dislocations after Hip Joint Replacement Surgery in Medan. This research was conducted at Haji Adam Malik General Hospital and several network hospitals of the Department of Orthopedics and Traumatology, USU Medical Faculty, located in Medan. This research was conducted from September 2021 to August 2022. There were a total of 32 samples included in this study, which were divided into 2 groups, namely cemented and uncemented who met the inclusion and exclusion criteria. The inclusion criteria for this study were all patients with closed neck femur fracture age more than 50 years old and exclusion criteria for this study are patients with fracture of both legs, osteomyelitis, malunion fracture, history of malignancy and currently undergoing chemotherapy treatment, sepsis, and total paralysis due to spinal cord injury or stroke.

3. Results

A total of 32 subjects were assessed for eligibility and met the inclusion criteria. Respondents were generally female (62.5%). All descriptive parameter data are presented in table 1.

Table 1. Basic Characteristics of Respondents

Characteristic	n	%
Sex		
Male	12	37.5%
Female	20	62.5%
Age (years)		
Rerata $\pm$ SD	72.21 $\pm$ 10.24	
Usia Termuda	54	
Usia Tertua	91	
Location		
Right	15	46.9%
Left	17	53.1%
Surgery Methods		
Bipolar Cemented	16	50%
Bipolar Uncemented	16	50%
Dislocation Post Operation		
Bipolar Cemented	0	
Bipolar Uncemented	0	

To analyze the clinical function assessment of MHHS from cemented bipolar hemiarthroplasty with uncemented bipolar hemiarthroplasty, a distribution test was conducted to determine the distribution of the data obtained using the Shapiro-Wilk test. The results of the normality test using Saphiro Wilk showed that the research data were normally distributed with p-values of 0.458 and 0.810 (>0.05) respectively.

Table 2. Normality test results Modified Harris Hip Score

Surgery Methods	P Value
Bipolar Hemiarthroplasty Cemented	0.458
Bipolar Hemiarthroplasty Uncemented	0.810

From the results of the data normality test, it was found that the research data were normally distributed. Because the study data were normally distributed, the statistical test used to identify the presence of an external relationship between clinical function modified harris hip score in femoral neck fractures was the Independent T-Test (unpaired T-Test).

Statistical analysis of the clinical outcome assessment of modified harris hip score from cemented bipolar hemiarthroplasty and uncemented bipolar hemiarthroplasty in femoral neck fractures showed that there was no significant difference in modified harris hip score between cemented bipolar hemiarthroplasty and uncemented bipolar hemiarthroplasty with a significance value of 0.935 (> 0.05).

Table 3. Differences modified harris hip score in cemented and uncemented bipolar hemiarthroplasty

		Mean	P value
Modified Harris Hip Score	Bipolar Hemiarthroplasty	79.00±5.40	0.935
	Cemented		
	Bipolar Hemiarthroplasty	78.81±7.43	
	Uncemented		

4. Discussion

The incidence of fractures in the hip joint, both intracapsular and extracapsular, increases with age because bone mass will gradually decrease with age, causing osteoporosis. found in women. This can be related to hormonal transitions, especially during menopause, several studies have also shown that the risk factor for falling is greater in women than men, losing muscle mass in women occurs more quickly than men, especially after menopause. Women are also more susceptible to psychological disorders such as depression and often use various drugs that can increase the risk of falling.

Other groupings were made based on the location of the femoral neck fracture and the type of surgery (Table 1) and from this data it was obtained that the distribution of the number of samples based on the location of the fracture of the right femoral neck was 15 (46.9%) and the location of the fracture of the left femoral neck was 17 (53.1. %). Based on the type of operation made consecutively proportional that is half of each group. Of the 32 subjects 16 underwent cemented bipolar hemiarthroplasty and 16 underwent uncemented bipolar hemiarthroplasty surgery and no dislocation was found in the group that underwent cemented or uncemented bipolar hemiarthroplasty surgery.

To be able to analyze the comparison of MHHS function assessments from bipolar cemented hemiarthroplasty and uncemented hemiarthroplasty measures, the step that must be passed is testing the distribution of the samples obtained. The normality test of the data in this study was carried out using the Shapiro Wilk test, this test was carried out because the number of subjects obtained in the study was less than 50 subjects.

The results of this study showed that there was no significant difference in the comparison of clinical function outcomes of modified Harris hip score from cemented bipolar hemiarthroplasty and uncemented bipolar hemiarthroplasty in femoral neck fractures ($p=0.935$).

In a study conducted by Gupta et al in 2019 in India, cemented bipolar hemiarthroplasty was performed on 9 patients and uncemented bipolar hemiarthroplasty in 10 patients with a diagnosis of femoral neck fracture, evaluated for 6 weeks, 12 weeks and 1 year. At 6 weeks postoperatively there was no significant difference, but at 1 year after surgery, the clinical outcome of MHHS in the cemented group was better than the uncemented group.

In a study conducted by Bashir et al in 2020 in India, a study was conducted on 20 men and 30 women with a diagnosis of femoral neck fracture aged 61-88 years using uncemented modular bipolar hemiarthroplasty with a posterior approach evaluated at 6-11 months with a mean MHHS 88.96 and no postoperative dislocation.

In a study conducted by Movrin Igor in Slovenia in 2018 of 135 patients over the age of 76 years with femoral neck fractures were divided into 2 groups treated with cemented bipolar hemiarthroplasty and uncemented bipolar hemiarthroplasty, the two groups of patients above were divided not based on certain guidelines but based on considerations separately from the orthopedic specialist. The clinical outcome of HHS in the cemented group 6 weeks postoperatively 77.1 ± 13.1 and uncemented 71.3 ± 16.3 , at 24 months postoperative cemented group 81.2 ± 9.5 and uncemented group 79.6 ± 8.4 .

References

- Bucholz, Robert W, Heckman James D, Court-Brown, Charles M. (2006). *Rockwood and Green's Fracture in Adults*. 6th edition. Philadelphia: Lippincott Williams & Wilkins.
- Blomfeldt, R., Törnkvist, H., Eriksson, K., Söderqvist, A., Ponzer, S. and Tidermark, J., (2007). A randomised controlled trial comparing bipolar hemiarthroplasty with total hip replacement for displaced intracapsular fractures of the femoral neck in elderly patients. *The Journal of Bone and Joint Surgery. British volume*, 89-B(2), pp.160-165.
- Hadisoebroto Dilogio, Ismail & Djaja, Yoshi. (2013). Hemi or Total Hip Arthroplasty in Hip Fractures: What is the Contemporary Recommendations?: Real Scenario and How I Select My Patients.
- Ringshawl, Z., Bashir, Z., Bashir, A., Farooq, M. and Maqbool Wani, M., (2020). Functional Outcome of Uncemented Modular Bipolar Hemiarthroplasty Using Modified Harris Hip Score for Fractures of Femoral Neck in Elderly Patients. *Journal of Medical Science And clinical Research*, 08(09).
- Daddimani R, Patil V, Shettar C. RESULTS OF FRACTURE NECK OF FEMUR TREATED WITH MODULAR BIPOLAR HEMIARTHROPLASTY IN ELDERLY-OUR EXPERIENCE IN 70 CASES. *International Journal of Orthopaedics Traumatology and Surgical Sciences*. 2016 July;2(1):197-202.
- Kho D, Kim K, Kim H. and Kim D. (2008). Bipolar Hemiarthroplasty Using Calcar Replacement Stem for Hip Fractures in the Elderly. *Journal of the Korean Fracture Society*, 21(3): 232.
- Drake R, Vogl W, Mitchell A, Tibbitts R, Richardson P and Horn A. (2016). *Gray's basic anatomy*. 2nd ed. Philadelphia: Elsevier: 266-306.
- Egol, K., Koval, K., & Zuckerman, J. (2020). *Handbook of fractures* (6th ed). Philadelphia: Wolters Kluwer. pp. 366-367
- Salminen, S., 2005. *FEMORAL SHAFT FRACTURES IN ADULTS: EPIDEMIOLOGY, FRACTURE PATTERNS, NONUNIONS, AND FATIGUE FRACTURES*. Helsinki: The Faculty of Medicine of the University of Helsinki.
- Jo Mark J, Tencer Allan F, Gardner Michael J. (2015). *Rockwood and Green's fractures in adults*. 8th ed. Biomechanics of Fracture and Fracture Fixation Philadelphia: Wolters Kluwer. p42
- Mittal, R., & Banerjee, S. (2012). Proximal femoral fractures: Principles of management and review of literature. *Journal Of Clinical Orthopaedics And Trauma*, 3(1), 15-23. doi: 10.1016/j.jcot.2012.04.001
- Kain Michael S, Tortetta III Paul. (2015). Hip Dislocation and Fractures of The Femoral Head. *Rockwood and Green's fractures in adults*. 8th ed. Biomechanics of Fracture and Fracture Fixation Philadelphia: Wolters Kluwer. p1983-2030
- Keating John F. (2015). Femoral Neck Fracture. *Rockwood and Green's fractures in adults*. 8th ed. Biomechanics of Fracture and Fracture Fixation Philadelphia: Wolters Kluwer. p1983-2030
- Vanrusselt, J., Vansevenant, M., Vanderschueren, G., & Vanhoenacker, F. (2015). Postoperative radiograph of the hip arthroplasty: what the radiologist should know. *Insights Into Imaging*, 6(6), 591-600. doi: 10.1007/s13244-015-0438-5
- Mercer, W., & Sivananthan, S. (2012). *Mercer's textbook of orthopaedics and trauma*. 10th ed. London: Hodder Arnold. p. 343

- Azar, F., Beaty, J., Canale, S., & Campbell, W. (2017). *Campbell's operative orthopaedics*. 13th ed. Philadelphia: Elsevier. pp. 171-219
- Raaymakers E, Schipper I, Simmermacher R. Posterolateral approach to the hip. (2010). Retrieved from <https://surgeryreference.aofoundation.org/orthopedic-trauma/adult-trauma/proximal-femur/approach/posterolateral-approach-to-the-hip>
- Raaymakers E, Schipper I, Simmermacher R. Anterolateral approach for ORIF (femoral neck fractures), possible for prosthesis (neck or head fractures). (2010). Retrieved from <https://surgeryreference.aofoundation.org/orthopedic-trauma/adult-trauma/proximal-femur/approach/posterolateral-approach-to-the-hip>
- Raaymakers E, Schipper I, Simmermacher R. Direct lateral approach. (2010). Retrieved from <https://surgeryreference.aofoundation.org/orthopedic-trauma/adult-trauma/proximal-femur/approach/posterolateral-approach-to-the-hip>
- Donegan D, Huo M, Leslie M. Direct anterior approach. (2010). Retrieved from <https://surgeryreference.aofoundation.org/orthopedic-trauma/adult-trauma/proximal-femur/approach/posterolateral-approach-to-the-hip>
- Edwards, P., Queen, R., Butler, R., Bolognesi, M., & Lowry Barnes, C. (2016). Are Range of Motion Measurements Needed When Calculating the Harris Hip Score?. *The Journal Of Arthroplasty*, 31(4), 815-819. doi: 10.1016/j.arth.2015.10.016
- Soderman, P., & Malchau, H. (2001). Is the Harris Hip Score System Useful to Study the Outcome of Total Hip Replacement?. *Clinical Orthopaedics And Related Research*, 384, 189-197. doi: 10.1097/00003086-200103000-00022
- Sharma, S., Shah, R., Draviraj, K., & Bhamra, M. (2005). Use of telephone interviews to follow up patients after total hip replacement. *Journal Of Telemedicine And Telecare*, 11(4), 211-214. doi: 10.1258/1357633054068883
- Kumar, P., Sen, R., Aggarwal, S., Agarwal, S., & Rajnish, R. (2019). Reliability of Modified Harris Hip Score as a tool for outcome evaluation of Total Hip Replacements in Indian population. *Journal Of Clinical Orthopaedics And Trauma*, 10(1), 128-130. doi: 10.1016/j.jcot.2017.11.01
- Jones, C., Briffa, N., Jacob, J., & Hargrove, R. (2017). The Dislocated Hip Hemiarthroplasty: Current Concepts of Etiological factors and Management. *The Open Orthopaedics Journal*, 11(1), 1200-1212. doi: 10.2174/1874325001711011200
- Figved, W., Opland, V., Frihagen, F., Jervidalo, T., Madsen, J., & Nordsletten, L. (2009). Cemented versus Uncemented Hemiarthroplasty for Displaced Femoral Neck Fractures. *Clinical Orthopaedics & Related Research*, 467(9), 2426-2435. doi: 10.1007/s11999-008-0672-y
- Langslet, E., Frihagen, F., Opland, V., Madsen, J., Nordsletten, L., & Figved, W. (2014). Cemented versus Uncemented Hemiarthroplasty for Displaced Femoral Neck Fractures: 5-year Followup of a Randomized Trial. *Clinical Orthopaedics & Related Research*, 472(4), 1291-1299. doi: 10.1007/s11999-013-3308-9
- Jobory, A., Kärrholm, J., Hansson, S., Åkesson, K., & Rogmark, C. (2021). Dislocation of hemiarthroplasty after hip fracture is common and the risk is increased with posterior approach: result from a national cohort of

- 25,678 individuals in the Swedish Hip Arthroplasty Register. *Acta Orthopaedica*, 92(4), 413-418. doi: 10.1080/17453674.2021.1906517
- Enocson, A., Tidermark, J., Törnkvist, H., & Lapidus, L. (2008). Dislocation of hemiarthroplasty after femoral neck fracture: Better outcome after the anterolateral approach in a prospective cohort study on 739 consecutive hips. *Acta Orthopaedica*, 79(2), 211-217. doi: 10.1080/17453670710014996
- Gupta, V., Kumar Tiwari, P., Galfat, D., & Khichar, H. (2019). MANAGEMENT OF FRACTURE NECK FEMUR WITH CEMENTED AND UNCEMENTED HEMIARTHROPLASTY IN ELDERLY PATIENTS- COMPARISON OF OUTCOME THEREOF. *National Journal Of Medical And Dental Research*, 7(3), 622-626.
- Kerakkanavar, S., Kaladagi, D., Sanakal, N., Kaladagi, P., & Hongal, P. (2021). Evaluation of functional outcome of displaced fracture neck of femur in elderly treated with cemented bipolar prosthesis by modified Harris hip score. *International Journal Of Research In Orthopaedics*, 7(3), 561. doi: 10.18203/issn.2455-4510.intjresorthop20211610
- Pfeilschifter, J., Cooper, C., Watts, N., Flahive, J., Saag, K., & Adachi, J. et al. (2011). Regional and age-related variations in the proportions of hip fractures and major fractures among postmenopausal women: the Global Longitudinal Study of Osteoporosis in Women. *Osteoporosis International*, 23(8), 2179-2188. doi: 10.1007/s00198-011-1840-6
- Talbot, L., Musiol, R., Witham, E., & Metter, E. (2005). Falls in young, middle-aged and older community dwelling adults: perceived cause, environmental factors and injury. *BMC Public Health*, 5(1). doi: 10.1186/1471-2458-5-86
- Hornbrook, M., Stevens, V., Wingfield, D., Hollis, J., Greenlick, M., & Ory, M. (1994). Preventing Falls Among Community-Dwelling Older Persons: Results From a Randomized Trial. *The Gerontologist*, 34(1), 16-23. doi: 10.1093/geront/34.1.16
- Bhise, S., Borkar, S., Desai, S., & Kanode, N. (2018). Comparative Study of Patients Who Underwent Hemiarthroplasty of Hip by Anterolateral and Posterolateral Approach. *International Journal Of Scientific Study*, 6(2).
- Movrin, I. (2020). Cemented versus uncemented hemiarthroplasty for displaced femoral neck fractures: A randomized controlled trial with two years follow-up. *Acta Orthopaedica Et Traumatologica Turcica*, 54(1), 83-88. doi: 10.5152/j.aott.2020.01.432
- Stronach, B., Bergin, P., Perez, J., Watson, S., Jones, L., McGwin, G., & Ponce, B. (2019). The rising use of total hip arthroplasty for femoral neck fractures in the United States. *HIP International*, 30(1), 107-113. doi: 10.1177/1120700019832989
- Marya, S., Thukral, R., Hasan, R., & Tripathi, M. (2011). Cementless bipolar hemiarthroplasty in femoral neck fractures in elderly. *Indian Journal Of Orthopaedics*, 45(3), 236-242. doi: 10.4103/0019-5413.80042
- Moon, N., Shin, W., Do, M., Kang, S., Lee, S., & Suh, K. (2021). High conversion rate to total hip arthroplasty after hemiarthroplasty in young patients with a minimum 10 years follow-up. *BMC Musculoskeletal Disorders*, 22(1). doi: 10.1186/s12891-021-04153-4