

Negative Correlation of Smoking Habits With Hemoglobin Levels and Maximum Oxygen Volume Capacity on Students

Imam Fauzi^a, Olivia Andiana^a, Slamet Raharjo^{a*}

**Corresponding author: Slamet Raharjo, E-mail: slamet.raharjo.fik@um.ac.id*

a. Department of Sport Science, Faculty of Sport Science, Universitas Negeri Malang, Semarang No. 5 Street, Malang, East Java 65145, Indonesia

Abstract

Smoking habits can potentially affect the decrease in hemoglobin levels and VO₂max capacity. The purpose of this study is to investigate the association between smoking behaviors and hemoglobin levels as well as VO₂max capacity in college students. This is an analytic research design with a cross-sectional study. In total, there were 14 students had an average age of 21.93 ± 1.38 years, with a normal body mass index of 21.53 ± 1.36 kg/m², and an average smoking habit of 1881.07 ± 1907.01 cigarettes/year registered as research subjects. Data collection on smoking habits was carried out using interview survey techniques, hemoglobin levels were measured using Easy Touch GCHb with units of g/dL levels, and the multi-stage 20-m Shuttle Run Fitness Test was used to determine maximum oxygen volume (VO₂max). The Pearson product-moment correlation coefficient test with a significance threshold of 5% is used in the data analysis approach. The findings revealed a detrimental relationship between smoking behaviors and hemoglobin levels ($r=-0.834$; $p=0.000$) and VO₂max capacity ($r=-0.669$; $p=0.009$). The study's findings show that smoking behaviors have a detrimental association with hemoglobin levels and VO₂max capacity. This shows that the higher the cigarette consumption, the lower the hemoglobin level and VO₂max capacity.

Keywords: Smoking habit, hemoglobin level, VO₂max

1. Introduction

Smoking is the activity of smoking processed tobacco wrapped in white paper that is burned (Pradhana & Rochmania, 2018). In theory, the factor contributing to smoking habits is the development of nicotine addiction cigarettes by experiencing the beneficial effects of nicotine, namely for relaxation or calm, and reducing anxiety or tension (Fikriyah & Febrijanto, 2012). Smoking activity in Indonesia on average begins at the age of 15 years (Salsabila et al., 2022). At a young age in East Java, as many as 68.68% of smokers spend at least 1-5 cigarettes a day (Almaidah et al., 2021). Most smokers smoke at an interval of ≥ 60 minutes from waking up and cannot bear not to smoke in prohibited places (Salsabila et al., 2022).

The negative consequences of smoking, particularly on the respiratory organs (Rizqi & Rochmania, 2018). Smoking significantly reduces the function of the cardiorespiratory system by reducing the body's capacity to transport oxygen (de Borba et al., 2014). In addition, the main harmful substance that enters is carbon monoxide (CO) gas which worsens hemoglobin (Hb) and oxygen (O₂) levels (Putra et al., 2018), so smoking habits can reduce the quality of physical endurance (Febriadi & Andrijanto, 2021). However, little has been reported on the relationship between smoking habits and hemoglobin levels, and maximal oxygen volume.

Therefore, the purpose of this study is to examine the association between smoking behaviors and hemoglobin levels and maximal oxygen volume ($VO_2\text{max}$) in college students.

2. Material and Methods

This was a cross-sectional study with an analytic research approach. 14 students in total had an average age of 21.93 ± 1.38 years, with a normal body mass index of $21.53 \pm 1.36 \text{ kg/m}^2$, and had an average smoking habit of 1881.07 ± 1907.01 cigarettes/year registered as research subjects. Data collection on smoking habits was carried out using an interview survey technique (Amelia et al., 2016), hemoglobin levels were measured using Easy Touch GCHb (Easy Touch, Hsinchu, Taiwan) with g/dL levels (Raharjo et al., 2021), and The multi-stage 20-m Shuttle Run Fitness Test was used to determine maximum oxygen volume ($VO_2\text{max}$). (Puspodari et al., 2022). Data collection on smoking habits, hemoglobin levels, and $VO_2\text{max}$ was carried out at one time. The Pearson product-moment correlation coefficient test with a significance threshold of 5% is used in the data analysis approach.

3. Results

The results of the study reported that the subjects had an average age of 21.93 ± 1.38 years, with a normal body mass index of $21.53 \pm 1.36 \text{ kg/m}^2$. The subjects' average smoking habits were 1881.07 ± 1907.01 cigarettes/year, average hemoglobin $15.34 \pm 1.23 \text{ g/dL}$, average $VO_2\text{max}$ $28.14 \pm 4.53 \text{ mL/kg/min}$, and more details can be seen in Table 1 below. Meanwhile, the results of the correlation analysis between smoking habits and hemoglobin levels and $VO_2\text{max}$ show a negative correlation between the three which can be seen in Table 2.

Table 1. Characteristics of research subjects

Parameters	n	Minimum	Maximum	Mean	Std. Deviation
Age (yrs)	14	19.00	25.00	21.93	1.38
Height (m)	14	1.60	1.74	1.67	0.04
Weight (kg)	14	50.00	71.00	60.07	5.89
Body mass index (kg/m^2)	14	19.10	23.50	21.53	1.36
Smoking habits (cigarettes/year)	14	110.00	6570.00	1881.07	1907.01
Hemoglobin (g/dL)	14	12.20	17.00	15.34	1.23
$VO_2\text{max}$ (mL/kg/min)	14	23.90	40.00	28.14	4.53

Table 2. The results of the correlation analysis of smoking habits with hemoglobin and $VO_2\text{max}$ levels

Parameter	n	Smoking habits (Stems/yrs)	
		Pearson Correlation	Sig. (2-tailed)
Hemoglobin (g/dL)	14	-0.834**	0.000
$VO_2\text{max}$ (mL/kg/min)	14	-0.669**	0.009

4. Discussion

The mean hemoglobin level of the 14 research subjects was 15.34 g/dL , with the lowest value being 12.20 g/dL and the highest value being 17.00 g/dL . The correlation analysis results revealed a negative association between smoking practices and hemoglobin levels (Table 2). This result is different from the

research report by Amelia et al. (2016) indicated that there was no link between the Brinkman Index and the degree of smoking and hemoglobin levels. Many other factors affect hemoglobin levels in the body, especially for smokers (Septiani, 2022). Smoking is known to be a major public health problem and is considered the leading cause of preventable death worldwide (Wang et al., 2023). Results by Vivek et al. (2023) confirmed that smoking tobacco has a strong relationship with iron deficiency anemia. Heavy metals and several persistent organic pollutants included in cigarettes can cause illnesses, such as cardiovascular disease (CVD), cancer, and chronic obstructive pulmonary disease (COPD) (Kim et al., 2023; Achanzar et al., 2001; Qin et al., 2009; Donaldson et al., 2010). However, it is well known that blood cells are produced in the bone marrow (Remley et al., 2023). The possible negative effects of smoking can have a direct impact on the quality of circulating blood cells. This might explain why there is a link between smoking behaviors and hemoglobin levels in this study. Therefore, future studies may compare smoking behaviors to bone marrow quality to identify the relationship between smoking habits and hemoglobin levels, yielding more precise results.

The findings revealed that there was a negative relationship between smoking habits and $VO_2\text{max}$ capacity. That is, someone who has a smoking habit tends to have a lower $VO_2\text{max}$ capacity compared to a non-smoker (Zuhdi & Yuliasrid, 2017). Mitochondrial respiration is an important component of cell metabolism (Smolina et al., 2023). Mitochondrial respiration can be represented by $VO_2\text{max}$ capacity. $VO_2\text{max}$ measures an organism's ability to incorporate oxygen into metabolism to get energy (Rosado-Pérez & Mendoza-Núñez et al., 2018). According to the findings of this study, there is a negative relationship between smoking behaviors and obesity and $VO_2\text{max}$ capacity. The possible mechanism linking smoking to a reduction in $VO_2\text{max}$ might be oxidative damage and inflammation caused by smoking. Smoking-induced oxidative stress and inflammation are linked to the pathological processes of several chronic respiratory disorders, including asthma, emphysema, chronic obstructive pulmonary disease, and cancer (Xue et al., 2023). According to Decker et al. (2023), cigarette smoke directly impairs the mitochondrial respiration of carbohydrate-derived substrates, which is the main mechanism underlying cigarette smoke-induced muscle dysfunction, resulting in adverse effects such as excess glucose conversion to fatty acids and lipotoxicity. It is well understood that mitochondria play a crucial role in the regulation of the inflammatory response (Marchi et al., 2023).

5. Conclusion

The study's findings demonstrate that there was a negative relationship between smoking habits and hemoglobin levels and $VO_2\text{max}$ capacity. This shows that the higher the cigarette consumption, the lower the hemoglobin level and $VO_2\text{max}$ capacity.

References

- Achanzar, W. E., Diwan, B. A., Liu, J., Quader, S. T., Webber, M. M., & Waalkes, M. P. (2001). Cadmium-induced malignant transformation of human prostate epithelial cells. *Cancer Research*, 61(2), 455–458.
- Almaidah, F., Khairunnisa, S., Sari, I. P., Chrisna, C. D., Firdaus, A., Kamiliya, Z. H., & Puspitasari, H. P. (2021). Survey of Factors Causing Teen Smokers to Maintain Smoking Behavior. *Journal of Community Pharmacy*, 8(1), 20–26.
- Amelia, R., Nasrul, E., & Basyar, M. (2016). Correlation between degree of smoking based on Brinkman index and hemoglobin level. *Jurnal Kesehatan Andalas*, 5(3), 619–624.
- Decker, S. T., Matias, A. A., Bannon, S. T., Madden, J. P., Alexandrou-Majaj, N., & Layec, G. (2023). Effects of cigarette smoke on in situ mitochondrial substrate oxidation of slow- and fast-twitch skeletal muscles. *Life sciences*, 315, 121376. <https://doi.org/10.1016/j.lfs.2023.121376>.
- de Borja, A. T., Jost, R. T., Gass, R., Nedel, F. B., Cardoso, D. M., Pohl, H. H., Reckziegel, M. B., Corbellini, V. A., & Paiva, D. N. (2014). The influence of active and passive smoking on the cardiorespiratory fitness of adults. *Multidisciplinary respiratory medicine*, 9(1), 34. <https://doi.org/10.1186/2049-6958-9-34>.

- Donaldson, S. G., Van Oostdam, J., Tikhonov, C., Feeley, M., Armstrong, B., Ayotte, P., Boucher, O., Bowers, W., Chan, L., Dallaire, F., Dallaire, R., Dewailly, E., Edwards, J., Egeland, G. M., Fontaine, J., Furgal, C., Leech, T., Loring, E., Muckle, G., Nancarrow, T., et al. (2010). Environmental contaminants and human health in the Canadian Arctic. *The Science of the total environment*, 408(22), 5165–5234. <https://doi.org/10.1016/j.scitotenv.2010.04.059>.
- Febriadi, S. G. P., & Andrijanto, D. Comparison of Physical Fitness in Smokers and Non-Smokers Students at SMA Negeri 1 Porong. *Journal of Sports and Health Education*, 9(1), 229-232.
- Fikriyah, S., & Febrijanto, Y. (2012). Factors influencing smoking behavior in male students in the male dormitory. *Stikes journal*, 5(1), 99-109.
- Kim, J., Song, H., Lee, J., Kim, Y. J., Chung, H. S., Yu, J. M., Jang, G., Park, R., Chung, W., Oh, C. M., & Moon, S. (2023). Smoking and passive smoking increases mortality through the mediation effect of cadmium exposure in the United States. *Scientific reports*, 13(1), 3878. <https://doi.org/10.1038/s41598-023-30988-z>.
- Marchi, S., Guilbaud, E., Tait, S. W. G., Yamazaki, T., & Galluzzi, L. (2023). Mitochondrial control of inflammation. *Nature reviews. Immunology*, 23(3), 159–173. <https://doi.org/10.1038/s41577-022-00760-x>.
- Pradhana, A. S. G., & Rochmania, A. (2018). Analysis of the Physical Conditions of Athletes who smoke and do not smoke (Study of Men's Volleyball Athletes in Nanggala Surabaya, Age 15-16, 2018). *Sports Achievement Journal*, 1(2). 1-5.
- Puspodari, P., Wiriawan, O., Setijono, H., Arfanda, P. E., Himawanto, W., Koestanto, S. H., Hantoro, B., Lusianti, S., Putra, R. P., Prasetyo, R., & Pranoto, A. (2022). Effectiveness of Zumba Exercise on Maximum Oxygen Volume, Agility, and Muscle Power in Female Students. *Physical Education Theory and Methodology*, 22(4), 478–484. <https://doi.org/10.17309/tmf.2022.4.04>.
- Putra, Y. V. C. (2018). Survey of VO2 Max Smoker and Non-Smoker Levels in High School Students. *ACTIVE: Journal of Physical Education, Sport, Health and Recreation*, 7(3), 120-123.
- Qin, Y. Y., Leung, C. K. M., Leung, A. O. W., Wu, S. C., Zheng, J. S., & Wong, M. H. (2009). Persistent organic pollutants and heavy metals in adipose tissues of patients with uterine leiomyomas and the association of these pollutants with seafood diet, BMI, and age. In *Environmental Science and Pollution Research* (Vol. 17, Issue 1, pp. 229–240). Springer Science and Business Media LLC. <https://doi.org/10.1007/s11356-009-0251-0>.
- Raharjo, S., Pranoto, A., Rejeki, P. S., Harisman, A. S. M., Pamungkas, Y. P., & Andiana, O. (2021). Negative Correlation between Serum Brain-derived Neurotrophic Factor Levels and Obesity Predictor Markers and Inflammation Levels in Females with Obesity. *Open Access Macedonian Journal of Medical Sciences*, 9(B), 1021–1026. <https://doi.org/10.3889/oamjms.2021.6840>.
- Remley, V. A., Collins, A., Underwood, S., Jin, J., Kim, Y., Cai, Y., Prochazkova, M., Moses, L., Byrne, K. M., Jin, P., Stroneck, D. F., & Highfill, S. L. (2023). Optimizing a fully automated and closed system process for red blood cell reduction of human bone marrow products. *Cytotherapy*, 25(4), 442–450. <https://doi.org/10.1016/j.jcyt.2022.12.006>.
- Rizqi, A., & Rochmania, A. (2020). Comparison of Maximum Oxygen Volume (VO₂max) Athletes Smokers and Non-Smokers (Study on the Asmara FC U18 Boys Futsal Team, Pamekasan Regency). *Sports Achievement Journal*, 3(3).
- Rosado-Pérez, J., & Mendoza-Núñez, V. M. (2018). Relationship Between Aerobic Capacity With Oxidative Stress and Inflammation Biomarkers in the Blood of Older Mexican Urban-Dwelling Population. Dose-response: a publication of International Hormesis Society, 16(2), 1559325818773000. <https://doi.org/10.1177/1559325818773000>.
- Salsabila, N.N., Indraswari, N., & Sujatmiko, B. (2022). Description of smoking habits in Indonesia based on Indonesia Family Life Survey 5 (IFLS 5). *Journal of Indonesian Health Economics*, 7(1), 13-22.
- Septiani, R. (2022). Relationship between smoking duration and smoking frequency with hemoglobin (Hb) levels in active smokers. *Babul Ilmi Scientific Journal of Health Multi-Science*, 14(1), 30-40.
- Smolina, N., Khudiakov, A., & Kostareva, A. (2023). Assaying Mitochondrial Respiration as an Indicator of Cellular Metabolism and Fitness. *Methods in molecular biology* (Clifton, N.J.), 2644, 3–14. https://doi.org/10.1007/978-1-0716-3052-5_1.
- Xue, J., Li, Z., Li, X., Hua, C., Shang, P., Zhao, J., Liu, K., & Xie, F. (2023). Evaluation of cigarette smoke-induced oxidative stress and inflammation in BEAS-2B cells based on a lung microfluidic chip. *Food and chemical toxicology: an international journal published for the British Industrial Biological Research Association*, 176, 113787. <https://doi.org/10.1016/j.fct.2023.113787>.
- Vivek, A., Kaushik, R. M., & Kaushik, R. (2023). Tobacco smoking-related risk for iron deficiency anemia: A case-control study. *Journal of addictive diseases*, 41(2), 128–136. <https://doi.org/10.1080/10550887.2022.2080627>.
- Wang, J., Wang, Y., Zhou, W., Huang, Y., & Yang, J. (2023). Impacts of cigarette smoking on blood circulation: do we need a new approach to blood donor selection? *Journal of health, population, and nutrition*, 42(1), 62. <https://doi.org/10.1186/s41043-023-00405-2>.
- Zuhdi, A. J. (2016). The Relationship between Smoking Habits and Maximum Oxygen Volume (VO₂max) in Students of the 2015 Unesa Department of Health and Research. *Journal of Sports Health*, 4(4), 81-89.