

Effects of High-Intensity Interval Exercise on Interleukin 6 Levels in Healthy Male Adolescents

Olivia Andiana^a, Mahmud Yunus^a, Slamet Raharjo^{a,*}, Ulul Azmi AnnurPutra^a,
 Raja Mohammed Firhad Raja Azidin^b

^{*}*slamet.raharjo.fik@um.ac.id*

^a*Department of Sport Science, Faculty of Sport Science, Universitas Negeri Malang, Semarang No. 5, Malang, East Java 65145, Indonesia*

^b*Faculty of Sports Science and Recreation, Universiti Teknologi MARA, 40450 Shah Alam, Malaysia*

Abstract

This study aims to prove the effects of high-intensity interval exercise on interleukin 6 (IL-6) levels in healthy male adolescents. Twenty healthy male adolescents aged twenty to twenty-five years with a body mass index (BMI) of 19-24 kg/m² were given one session of high-intensity interval exercise intervention by pedaling an ergocycle for 40 minutes. The ELISA method evaluated IL-6 levels between pretest and posttest in both groups. The data analysis technique uses a Paired Sample T-Test with a significance level of 5%. The results showed that there was an increase in the average IL-6 levels between the pretest and posttest in the high-intensity interval exercise group ($p \leq 0.05$). In contrast, in the control group there was no significant difference in the average IL-6 levels. 6 between pretest and posttest ($p \geq 0.05$). It is proven that one session of high-intensity interval exercise increases IL-6 levels in healthy male adolescents.

Keywords: High-intensity interval exercise; ergocycle exercise; IL-6 levels; metabolism

1. Introduction

Interleukin 6 (IL-6) is a cytokine that has pleiotropic functions in various tissues and organs (Muñoz-Cánoves et al., 2013). Skeletal muscle can produce and release IL-6 significantly after exercise (Muñoz-Cánoves et al., 2013; Lee & Jun, 2019). The increase in IL-6 after exercise depends on several factors such as age, body composition, and genetically expressed variations in the inflammatory response, including greater expression of toll-like receptor 4 (TLR4), which can induce greater transcription of inflammatory cytokines (Rogerio & Calder, 2018). When released into the circulation, IL-6 can promote metabolic modulation in several organs like hormones, such as hepatic glucose production during exercise or lipolysis occurring in adipose tissue (Thyfault & Bergouignan, 2020). In skeletal muscle, IL-6 will also aid glucose metabolism in a manner dependent on signal transduction involving PI-3 kinase and the JAK-STAT pathway through enhanced expression of the major glycolytic enzymes, hexokinase 2 and PFKFB-3 (Kumari et al., 2016). This mechanism will also increase glucose transport by inducing the expression of glucose transporters 1 and 4

(GLUT-1 and GLUT-4), then translocating to the plasma membrane to increase glucose absorption (Kumari et al., 2016). In addition, IL-6 also causes activation of AMP-activated protein kinase (AMPK) phosphorylation, which is useful for increasing glucose uptake and lipid oxidation (Carey et al., 2006). The acute release of IL-6 after exercise can also produce anti-inflammatory effects characterized by increased production of anti-inflammatory cytokines, such as IL-1ra and IL-10 (Tajra et al., 2014).

Acute physical exercise has a positive effect in inhibiting the development of chronic diseases through post-exercise inflammatory responses (Pranoto et al., 2023; Kramer & Goodyear, 2007). One of the inflammatory responses that increases post-exercise is IL-6 levels (Fischer, 2006; Pedersen & Febbraio, 2008). IL-6 levels increase exponentially (up to 100-fold) in response to exercise and decrease rapidly in the post-exercise period (Docherty et al., 2022). The magnitude of the increase in IL-6 levels during acute exercise depends on the training mode, training intensity, training duration, and training capacity of each individual (Ostrowski et al., 1999; Ostrowski et al., 2000; Febbraio & Pedersen, 2002). Several studies reported inconsistent results (increase, decrease, or no change) in IL-6 levels after acute exercise (Dimitrov et al., 2017; Windsor et al., 2018; Mendham et al., 2011). Factors that are likely to influence the inconsistency of results on IL-6 levels, namely differences in training mode, training intensity, and training duration (Mendham et al., 2011; Cerqueira et al., 2020; Pranoto et al., 2023).

Based on this, this research aims to prove the effects of high-intensity interval exercise on interleukin 6 (IL-6) levels in healthy male adolescents.

2. Material and methods

This research uses a true-experimental method with a pretest-posttest control group design research design. Respondents in this study were twenty healthy male adolescents, aged twenty to twenty-five years, with body mass index (BMI) of 19-24 kg/m², normal blood pressure, normal heart rate, normal fasting blood glucose, and normal hemoglobin. Respondents were randomly divided into two groups, namely CTL (n = 10, control group), and HIE (n = 10, high-intensity interval exercise group).

High-intensity interval exercise is done by pedaling an ergocycle for 40 minutes with details of 5 minutes of warm-up (50% – 60% HRmax), 30 minutes of core (80 – 90% HRmax) which is done in intervals (6 sets) each set is done for 5 minutes with active rest between sets for 2 minutes and closed with 5 minutes of cooling down (50% – 60% HRmax). The high-intensity interval exercise intervention is applied once (acute exercise). Monitor heart rate during exercise using Polar H7.

Blood was drawn from the cubital vein for 3 ml. When taking blood, the subject was in a sleeping position. Blood sampling was carried out 2 times, namely 30 minutes before the intervention and 10 minutes after the intervention. IL-6 levels were measured using the Enzyme Link Immunosorbent Assay (ELISA) method (Cat.No.:E-EL-H6156; Elabscience, Inc., USA).

The data analysis technique used SPSS version 21 software. The Shapiro-Wilk test and Levene test were used to test normality and homogeneity with a significant level ($p \geq 0.05$). The difference test uses a Paired Sample T-Test and an Independent Samples T-Test with a significant level ($p \leq 0.05$). Data presentation is done as mean $\pm$ SD.

3. Results

The differences in the results of the analysis of the characteristics of research subjects in each group can be seen in Table 1.

Table 1. Differences in characteristics of research subjects in the two groups

Variable	n	Groups		Sig. (2-tailed)
		CTL	HIE	
Age, years	10	20.60 (1.17)	21.20 (1.03)	0.241
Weight, kg	10	1.65 (0.05)	1.68 (0.04)	0.221
Height, m	10	62.01 (5.66)	63.22 (6.40)	0.660
BMI, kg/m ²	10	22.74 (1.21)	22.50 (1.76)	0.729
SBP, mmHg	10	116.60 (7.91)	117.70 (5.74)	0.727
DBP, mmHg	10	72.10 (7.16)	68.40 (3.98)	0.175
HR, bpm	10	71.90 (10.39)	70.80 (9.705)	0.810
SpO ₂ , %	10	98.00 (1.06)	97.20 (1.23)	0.136
BT, °C	10	35.26 (0.56)	35.45 (0.83)	0.557

Description: BMI: Body mass index; SBP: Systolic blood pressure; DBP: Diastolic blood pressure; HR: Heart rate; SpO₂: Oxygen saturation; BT: Body temperature. Sig. (2-tailed) was obtained using the Independent Sample T-Test. Data are presented as mean±SD.

Based on the results of statistical analysis, it shows that there is no significant difference in the average data on the characteristics of research subjects in each group ($p \leq 0.05$). The results of the analysis of differences in IL-6 levels between the pretest and posttest in each group are presented in Figure 1.




Figure 1. Differences in IL-6 level analysis results between pretest and posttest.

Description: Sig. (2-tailed) was obtained using the Independent Sample T-Test. Data are displayed as Mean±SD. (*) Significantly different from CTL ($p \leq 0.05$).

4. Discussion

Based on the research results, shows that high-intensity interval exercise (HIE) significantly increases IL-6 levels. These results are in line with the results of research conducted by Zwetsloot et al. (2014) who reported that a single session of high-intensity interval training (HIIT) induced an increase in IL-6 levels compared to pre-exercise in healthy young men. Another study reported that serum IL-6 levels in the HIIT group increased significantly in the one-hour post-HIIT phase in middle-aged men (Rohnejad & Monazzami, 2023). Exercise has complex benefits for the body (Rueggsegger & Booth, 2018). This is because exercise not only involves skeletal muscle contractions but also systemic activity of the respiratory and circulatory systems (Nara & Watanabe, 2021). During exercise, there is an increase in muscle contractions which will stimulate myocytes to produce myokines which are involved in the autocrine regulation of metabolism in muscles as well as in the paracrine/endocrine regulation of other tissues and organs including adipose tissue, liver, and brain through their receptors (Lee & Jun, 2019; Carson, 2017). This is one of the pleiotropic functions of IL-6 in various tissues and organs (Muñoz-Cánoves et al., 2013). Muscle movement during exercise creates stressors and secretes myokines, such as IL-6 (Lira et al., 2009). Increased IL-6 triggers the release of anti-inflammatory cytokines (IL-10 and IL-1R) which will balance the inflammatory events after exercise (Abd El-Kader & Al-Shreef, 2018). Muscle contraction will trigger the activation of hypothalamic–pituitary–adrenal axis (HPA-axis) to stimulate an increase in cortisol levels (Bonato et al., 2017), and increase potent anti-inflammatory hormones in the blood (Ruijters et al., 2016). Muscle contractions also trigger activation of the sympathetic nervous system to increase catecholamine levels. Catecholamines can suppress the production of lipopolysaccharide (LPS) which produces TNF- α and IL-1 β (Gleeson et al., 2011), thereby reducing levels of the pro-inflammatory cytokine, TNF- α to maintain the balance of inflammation in the body (Cerqueira et al., 2020).

IL-6 is known to be the first myokine discovered and is one of the most popular myokines discussed in several previous studies (Zunner et al., 2022; Mahmoud et al., 2022; Mageriu et al., 2020; Gonzalez-Gil & Elizondo-Montemayor, 2020; Furuichi et al., 2018; Gmiat et al., 2017; Karstoft Pedersen, 2016). Although it is known that IL-6 is a pro-inflammatory cytokine, it also has anti-inflammatory capacity when released by contracting muscles (Zunner et al., 2022). IL-6 has regenerative and anti-inflammatory functions, especially when secreted by skeletal muscles during physical exercise (Kistner et al., 2022). As an anti-inflammatory cytokine, IL-6 has been reported to inhibit pro-inflammatory effects, such as TNF- α , promote M2 macrophage polarization, and improve insulin sensitivity (Sindhu et al., 2015). Increases in IL-6 levels mediated by physical exercise can reduce the risk of various non-communicable chronic diseases, such as diabetes mellitus, dementia, cardiovascular disease, and some types of cancer (Severinsen & Pedersen, 2020). This proves that exercise has many health benefits, one of which is improving the body's metabolism (Thyfault & Bergouignan, 2020).

5. Conclusion

The results of the study prove that one session of high-intensity interval exercise (HIE) increases IL-6 levels in healthy male adolescents. Therefore, HIE can be used to improve and maintain the body's metabolism.

Acknowledgments

We thank the Faculty of Sports Science, Universitas Negeri Malang for the 2023 Decentralization of Research grant [No: 28.04.16/UN32.6.2/LT/2023].

References

- 1 Abd El-Kader, S. M., & Al-Shreef, F. M. (2018). Impact of aerobic exercises on selected inflammatory markers and immune system response among patients with sickle cell anemia in asymptomatic steady state. *African health sciences*, 18(1), 111–119. <https://doi.org/10.4314/ahs.v18i1.15>.
- 2 Bonato, M., La Torre, A., Saresella, M., Marventano, I., Merati, G., & Vitale, J. A. (2017). Salivary cortisol concentration after high-intensity interval exercise: Time of day and chronotype effect. *Chronobiology international*, 34(6), 698–707. <https://doi.org/10.1080/07420528.2017.1311336>.
- 3 Carey, A. L., Steinberg, G. R., Macaulay, S. L., Thomas, W. G., Holmes, A. G., Ramm, G., Prelovsek, O., Hohnen-Behrens, C., Watt, M. J., James, D. E., Kemp, B. E., Pedersen, B. K., & Febbraio, M. A. (2006). Interleukin-6 increases insulin-stimulated glucose disposal in humans and glucose uptake and fatty acid oxidation in vitro via AMP-activated protein kinase. *Diabetes*, 55(10), 2688–2697. <https://doi.org/10.2337/db05-1404>.
- 4 Carson B. P. (2017). The Potential Role of Contraction-Induced Myokines in the Regulation of Metabolic Function for the Prevention and Treatment of Type 2 Diabetes. *Frontiers in endocrinology*, 8, 97. <https://doi.org/10.3389/fendo.2017.00097>.
- 5 Cerqueira, É., Marinho, D. A., Neiva, H. P., & Lourenço, O. (2020). Inflammatory Effects of High and Moderate Intensity Exercise-A Systematic Review. *Frontiers in physiology*, 10, 1550. <https://doi.org/10.3389/fphys.2019.01550>.
- 6 Dimitrov, S., Hulteng, E., & Hong, S. (2017). Inflammation and exercise: Inhibition of monocytic intracellular TNF production by acute exercise via β 2-adrenergic activation. *Brain, behavior, and immunity*, 61, 60–68. <https://doi.org/10.1016/j.bbi.2016.12.017>.
- 7 Docherty, S., Harley, R., McAuley, J. J., Crowe, L., Pedret, C., Kirwan, P. D., Siebert, S., & Millar, N. L. (2022). The effect of exercise on cytokines: implications for musculoskeletal health: a narrative review. *BMC sports science, medicine & rehabilitation*, 14(1), 5. <https://doi.org/10.1186/s13102-022-00397-2>.
- 8 Febbraio, M. A., & Pedersen, B. K. (2002). Muscle-derived interleukin-6: mechanisms for activation and possible biological roles. *FASEB journal : official publication of the Federation of American Societies for Experimental Biology*, 16(11), 1335–1347. <https://doi.org/10.1096/fj.01-0876rev>.
- 9 Fischer C. P. (2006). Interleukin-6 in acute exercise and training: what is the biological relevance?. *Exercise immunology review*, 12, 6–33.
- 10 Furuichi, Y., Manabe, Y., Takagi, M., Aoki, M., & Fujii, N. L. (2018). Evidence for acute contraction-induced myokine secretion by C2C12 myotubes. *PloS one*, 13(10), e0206146. <https://doi.org/10.1371/journal.pone.0206146>.
- 11 Gleeson, M., Bishop, N. C., Stensel, D. J., Lindley, M. R., Mastana, S. S., & Nimmo, M. A. (2011). The anti-inflammatory effects of exercise: mechanisms and implications for the prevention and treatment of disease. *Nature reviews. Immunology*, 11(9), 607–615. <https://doi.org/10.1038/nri3041>.
- 12 Gmiat, A., Micielska, K., Kozłowska, M., Flis, D. J., Smaruj, M., Kujach, S., Jaworska, J., Lipińska, P., & Ziemann, E. (2017). The impact of a single bout of high intensity circuit training on myokines' concentrations and cognitive functions in women of different age. *Physiology & behavior*, 179, 290–297. <https://doi.org/10.1016/j.physbeh.2017.07.004>.

- 13 Gonzalez-Gil, A. M., & Elizondo-Montemayor, L. (2020). The Role of Exercise in the Interplay between Myokines, Hepatokines, Osteokines, Adipokines, and Modulation of Inflammation for Energy Substrate Redistribution and Fat Mass Loss: A Review. *Nutrients*, 12(6), 1899. <https://doi.org/10.3390/nu12061899>.
- 14 Karstoft, K., & Pedersen, B. K. (2016). Exercise and type 2 diabetes: focus on metabolism and inflammation. *Immunology and cell biology*, 94(2), 146–150. <https://doi.org/10.1038/icb.2015.101>.
- 15 Kistner, T. M., Pedersen, B. K., & Lieberman, D. E. (2022). Interleukin 6 as an energy allocator in muscle tissue. *Nature metabolism*, 4(2), 170–179. <https://doi.org/10.1038/s42255-022-00538-4>.
- 16 Kramer, H. F., & Goodyear, L. J. (2007). Exercise, MAPK, and NF-kappaB signaling in skeletal muscle. *Journal of applied physiology (Bethesda, Md. : 1985)*, 103(1), 388–395. <https://doi.org/10.1152/jappphysiol.00085.2007>.
- 17 Kumari, N., Dwarakanath, B. S., Das, A., & Bhatt, A. N. (2016). Role of interleukin-6 in cancer progression and therapeutic resistance. *Tumour biology : the journal of the International Society for Oncodevelopmental Biology and Medicine*, 37(9), 11553–11572. <https://doi.org/10.1007/s13277-016-5098-7>.
- 18 Lee, J. H., & Jun, H. S. (2019). Role of Myokines in Regulating Skeletal Muscle Mass and Function. *Frontiers in physiology*, 10, 42. <https://doi.org/10.3389/fphys.2019.00042>.
- 19 Lira, F. S., Koyama, C. H., Yamashita, A. S., Rosa, J. C., Zanchi, N. E., Batista, M. L., Jr, & Seelaender, M. C. (2009). Chronic exercise decreases cytokine production in healthy rat skeletal muscle. *Cell biochemistry and function*, 27(7), 458–461. <https://doi.org/10.1002/cbf.1594>.
- 20 Mageriu, V., Manole, E., Bastian, A. E., & Staniceanu, F. (2020). Role of Myokines in Myositis Pathogenesis and Their Potential to be New Therapeutic Targets in Idiopathic Inflammatory Myopathies. *Journal of immunology research*, 2020, 9079083. <https://doi.org/10.1155/2020/9079083>.
- 21 Mahmoud, N., Mohammadreza, H. A., Abdolhosein, T. K., Mehdi, N., & Arent, S. M. (2022). Serum myokine levels after linear and flexible non-linear periodized resistance training in overweight sedentary women. *European journal of sport science*, 22(4), 658–668. <https://doi.org/10.1080/17461391.2021.1895893>.
- 22 Mendham, A. E., Donges, C. E., Liberts, E. A., & Duffield, R. (2011). Effects of mode and intensity on the acute exercise-induced IL-6 and CRP responses in a sedentary, overweight population. *European journal of applied physiology*, 111(6), 1035–1045. <https://doi.org/10.1007/s00421-010-1724-z>.
- 23 Muñoz-Cánoves, P., Scheele, C., Pedersen, B. K., & Serrano, A. L. (2013). Interleukin-6 myokine signaling in skeletal muscle: a double-edged sword?. *The FEBS journal*, 280(17), 4131–4148. <https://doi.org/10.1111/febs.12338>.
- 24 Nara, H., & Watanabe, R. (2021). Anti-Inflammatory Effect of Muscle-Derived Interleukin-6 and Its Involvement in Lipid Metabolism. *International journal of molecular sciences*, 22(18), 9889. <https://doi.org/10.3390/ijms22189889>.
- 25 Ostrowski, K., Rohde, T., Asp, S., Schjerling, P., & Pedersen, B. K. (1999). Pro- and anti-inflammatory cytokine balance in strenuous exercise in humans. *The Journal of physiology*, 515 (Pt 1)(Pt 1), 287–291. <https://doi.org/10.1111/j.1469-7793.1999.287ad.x>.
- 26 Ostrowski, K., Schjerling, P., & Pedersen, B. K. (2000). Physical activity and plasma interleukin-6 in humans—effect of intensity of exercise. *European journal of applied physiology*, 83(6), 512–515. <https://doi.org/10.1007/s004210000312>.
- 27 Pedersen, B. K., & Febbraio, M. A. (2008). Muscle as an endocrine organ: focus on muscle-derived interleukin-6. *Physiological reviews*, 88(4), 1379–1406. <https://doi.org/10.1152/physrev.90100.2007>.
- 28 Pranoto, A., Cahyono, M. B. A., Yakobus, R., Izzatunnisa, N., Ramadhan, R. N., Rejeki, P. S., Miftahussurur, M., Effendi, W. I., Wungu, C. D. K., & Yamaoka, Y. (2023). Long-Term Resistance-Endurance Combined Training Reduces Pro-Inflammatory Cytokines in Young Adult Females with Obesity. *Sports (Basel, Switzerland)*, 11(3), 54. <https://doi.org/10.3390/sports11030054>.

- 29 Pranoto, A., Rejeki, P. S., Miftahussurur, M., Setiawan, H. K., Yosika, G. F., Munir, M., Maesaroh, S., Purwoto, S. P., Waritsu, C., & Yamaoka, Y. (2023). Single 30 min treadmill exercise session suppresses the production of pro-inflammatory cytokines and oxidative stress in obese female adolescents. *Journal of basic and clinical physiology and pharmacology*, 34(2), 235–242. <https://doi.org/10.1515/jbcpp-2022-0196>.
- 30 Rogero, M. M., & Calder, P. C. (2018). Obesity, Inflammation, Toll-Like Receptor 4 and Fatty Acids. *Nutrients*, 10(4), 432. <https://doi.org/10.3390/nu10040432>.
- 31 Rohnejad, B., & Monazzami, A. (2023). Effects of high-intensity intermittent training on some inflammatory and muscle damage indices in overweight middle-aged men. *Apunts Sports Medicine*, 58(217), 100404. <https://doi.org/10.1016/j.apunsm.2023.100404>.
- 32 Rueggsegger, G. N., & Booth, F. W. (2018). Health Benefits of Exercise. *Cold Spring Harbor perspectives in medicine*, 8(7), a029694. <https://doi.org/10.1101/cshperspect.a029694>.
- 33 Ruijters, E. J., Haenen, G. R., Willemsen, M., Weseler, A. R., & Bast, A. (2016). Food-Derived Bioactives Can Protect the Anti-Inflammatory Activity of Cortisol with Antioxidant-Dependent and -Independent Mechanisms. *International journal of molecular sciences*, 17(2), 239. <https://doi.org/10.3390/ijms17020239>.
- 34 Severinsen, M., & Pedersen, B. K. (2020). Muscle-Organ Crosstalk: The Emerging Roles of Myokines. *Endocrine reviews*, 41(4), 594–609. <https://doi.org/10.1210/edrev/bnaa016>.
- 35 Sindhu, S., Thomas, R., Shihab, P., Sriraman, D., Behbehani, K., & Ahmad, R. (2015). Obesity Is a Positive Modulator of IL-6R and IL-6 Expression in the Subcutaneous Adipose Tissue: Significance for Metabolic Inflammation. *PloS one*, 10(7), e0133494. <https://doi.org/10.1371/journal.pone.0133494>.
- 36 Tajra, V., Tibana, R. A., Vieira, D. C., de Farias, D. L., Teixeira, T. G., Funghetto, S. S., Silva, A. O., de Sousa, N. M., Willardson, J., Karnikowski, M. G., & Prestes, J. (2014). Identification of high responders for interleukin-6 and creatine kinase following acute eccentric resistance exercise in elderly obese women. *Journal of science and medicine in sport*, 17(6), 662–666. <https://doi.org/10.1016/j.jsams.2013.09.012>.
- 37 Thyfault, J. P., & Bergouignan, A. (2020). Exercise and metabolic health: beyond skeletal muscle. *Diabetologia*, 63(8), 1464–1474. <https://doi.org/10.1007/s00125-020-05177-6>.
- 38 Windsor, M. T., Bailey, T. G., Perissiou, M., Meital, L., Golledge, J., Russell, F. D., & Askew, C. D. (2018). Cytokine Responses to Acute Exercise in Healthy Older Adults: The Effect of Cardiorespiratory Fitness. *Frontiers in physiology*, 9, 203. <https://doi.org/10.3389/fphys.2018.00203>.
- 39 Zunner, B., Wachsmuth, N. B., Eckstein, M. L., Scherl, L., Schierbauer, J. R., Haupt, S., Stumpf, C., Reusch, L., & Moser, O. (2022). Myokines and Resistance Training: A Narrative Review. *International journal of molecular sciences*, 23(7), 3501. <https://doi.org/10.3390/ijms23073501>.
- 40 Zwetsloot, K. A., John, C. S., Lawrence, M. M., Battista, R. A., & Shanely, R. A. (2014). High-intensity interval training induces a modest systemic inflammatory response in active, young men. *Journal of inflammation research*, 7, 9–17. <https://doi.org/10.2147/JIR.S54721>.