

Herbal Medicine, Traditional Practice, and Nutraceutical as Prospective Alternative Therapy for Post COVID-19 Syndrome

Rahmi Nugraningrum^a, Maftuchah Rochmanti^{b*}

^b maftuchah-r@fk.unair.ac.id

^aMedical Study Program, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia, 60131

^bDepartment of Anatomy, Histology, and Pharmacology, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia, 60131

Abstract

Following COVID-19 infection, a post COVID-19 patient may experience Post COVID-19 Syndrome (PCS). This disorder is significantly important for therapies because it can lower an individual's quality of life following COVID-19 infection. Several treatment approaches, including non-conventional therapies such as herbal medicine, traditional therapy, and nutraceuticals, have been suggested for PCS. This narrative review will thoroughly deliberate the pathogenesis of PCS and several alternative, non-conventional therapy modalities, along with their efficacy and side effects. A literature search using PubMed and ScienceDirect databases, using the keywords “Post COVID-19” AND “Herbal Medicine” OR “nutraceutical” OR “traditional therapy” AND “efficacy” OR “safety” to find relevant studies. Clinical trials examining the effectiveness, safety, and indications of these new treatments are covered in our study. According to our research, these approaches show a lot of promise for being both safe and effective for treating post-COVID-19 syndrome. Therefore, more in-depth research is needed on some different therapy modalities that were not addressed in this study but may present interesting avenues for future post-COVID-19 syndrome treatment.

Keywords: herbal medicine; nutraceutical; post COVID-19 syndrome; infectious disease; traditional therapy

1. Introduction

In December 2019, a new coronavirus called SARS-CoV-2 (severe acute respiratory syndrome coronavirus) was discovered. Before recent discoveries, it was thought that the COVID-19 pandemic was mostly driven by human-to-human transmission and that it had first spread zoonotically through a seafood market in Wuhan, China (Zhang et al., 2020). By November 2021, it is expected that almost three billion people, or 44% of the world's population, will have at least one case of SARS-CoV-2 (Li et al., 2020). The symptoms of COVID-19 including fever, dyspnea, sore throat, chest discomfort, and neurological, cardiomyopathy, thromboembolism, and gastrointestinal disorders, are comparable to those of viral pneumonia (Hope and Evering, 2022). While some patients exhibit no symptoms at all, the majority of patients report mild to severe symptoms (Chen et al., 2022). While most treatments have been effective in assisting patients in making a full recovery from this infection, some of them continue to induce symptoms even after a COVID-19 test comes back negative. This condition called Post COVID-19 syndrome.

Post COVID-19 Syndrome (PCS) is a group of persistent symptoms that occur in who have had a history of suspected or proven SARS-CoV-2 infection. The symptoms usually start to show up three months after the COVID-19 infection started, last for at least two months and have no other known cause. Estimates from the WHO suggest that 200 million individuals are either experiencing or have previously had long-term health effects from COVID-19 (Chen et al., 2022). Ten percent to 35% of those recovering from acute illnesses are thought to be impacted by Post COVID-19 Syndrome; hospitalized patients are more likely to be afflicted (Hope and Evering, 2022). Several organs may be affected by PCS, which also has a remission-relapse propensity and a markedly reduced quality of life (Osmanov et al., 2022). Mast cell activation syndrome (MCAS) is linked to post-COVID-19 syndrome (PCS) and is typified by hyperactive mast cells releasing chemical mediators in excess and inappropriately. PCS can be caused by significant consequences from inflammatory reactions, host-microbiome alterations, reactivation of pathogens, and autoimmune mimicry (Jones et al., 2021).

To treat PCS, practitioners have tried their best. There are still large clinical and practical gaps in PCS treatment, even with the advent of several therapies. This does not provide suggestions for a specific therapy plan. Empirical data continues to be the cornerstone of therapy. Researchers are looking into how well histamine antagonist drugs like cimetidine, ranitidine, and famotidine, as well as antibiotics and antivirals like azithromycin, remdesivir, and favipiravir, work to suppress PCS. In the same way, monoclonal antibodies such as siltuximab, anakinra, and leronlimab work. Antidepressants may also enhance immune function and reduce peripheral inflammatory markers, according to some research (Koc et al., 2022). However the exact mechanism of action of these medications is unknown, and there are worries about long-term negative effects. Antivirals are one medication that can weaken the immune system. Patients become more vulnerable to opportunistic infections as a result.

Other treatment modalities, however, are also being researched since they are believed to be more effective and safe than conventional pharmacological therapy. A few instances of this are herbal medicine, and physical rehabilitation. This includes conventional methods such as tai chi, yoga, and acupuncture.

Herbal medication, such as Chinese Herbal medication (CHM), has been used in conjunction with conventional treatment to help prevent COVID-19 infection because of its various benefits (Chien et al., 2022). It has been demonstrated that Ayurveda and other commercial plant extracts can successfully alleviate fatigue and enhance overall health (Naureen et al., 2021). During the healing phase following an acute COVID-19 infection, non-pharmacological treatment management such as acupuncture, yoga, and Chinese martial arts (Tai Chi) is advised (Kim et al., 2022). The Acupuncture Association of China has even suggested guidelines for the use of moxibustion and acupuncture in treating COVID-19 and PCS (Zhang et al., 2020). Acupuncture, when paired with normal regimens, has been proven to be beneficial in relieving symptoms of anxiety, exhaustion, shortness of breath, loss of appetite, and insomnia with no negative side effects or consequences, according to several recent clinical trials (Jia et al., 2020).

Nutraceuticals, multivitamins, and mineral supplements, on the other hand, may be useful as PCS therapy possibilities because of their potential anti-inflammatory and anti-oxidative qualities (Singh et al., 2021). A study found that taking multivitamin supplements helped PCS patients' clinical symptoms (Koc et al., 2022; Rossato et al., 2021). Nicotinamide ribose, a kind of vitamin B3, is now showing promise in the treatment of chronic fatigue and cognitive impairment in two clinical trials. Probiotics, such as *Lactobacillus plantarum*, have been demonstrated in multiple studies to have beneficial effects on SARS-CoV-2-infected intestinal epithelial cells by functioning as an antiviral and by modifying the axis between the gut and the brain and the gut and lung (Liu et al., 2020).

The use of stem cells in treatment is gaining popularity among clinicians as well. In COVID-19 therapy, stem cells are commonly used as immunomodulatory agents to reduce the inflammatory process and prevent cytokine storms. Stem cells have been used in the past to treat a variety of diseases, such as autoimmune and viral disorders. Among the functions of stem cells include immunomodulation, regeneration potential, alveolar fluid clearance, and thrombotic prophylaxis (Jamshidi et al., 2021). The pathogenesis of PCS and potential alternative treatments, including herbal remedies, conventional therapy, and nutraceuticals, will be elaborated in detail in this paper.

2. Materials and Method

This review was conducted by searching for the efficacy, safety indication, and recommendation of using recent herbal medicine, traditional therapy, and nutraceutical for Post COVID-19 in two databases, PubMed (<https://pubmed.ncbi.nlm.nih.gov> and ScienceDirect(<https://www.sciencedirect.com/>). The Keyword is “Post COVID-19” AND “Herbal Medicine” OR “traditional therapy” OR “nutraceutical”, AND “Efficacy” OR “safety”.

The inclusion criteria are full-text literature and studies conducted within the last five years, and the research subjects were post COVID-19 patients who received herbal medicine, traditional therapy, or nutraceutical. The types of studies included in this review are in-silico studies, experimental (in vitro and in vivo), and clinical trials. The exclusion criteria were review articles without an English version.

3. Pathophysiology of Post COVID-19 Syndrome

The World Health Organization (WHO) states that the symptoms of post-COVID-19 syndrome (PCS) often appear three months after COVID-19 infection, persist for at least two months, and do not related with any other likely diagnosis. The two stages of PCS sequelae are (1) subacute or persistent COVID-19 symptoms, which include anomalies and manifestations and are identified by a negative Polymerase Chain Reaction (PCR) but the virus being found in the nasopharynx, that appear 4–12 weeks following the acute infection; and (2) Disorders and symptoms that manifest 12 weeks following the onset of the acute COVID-19 infection and are unrelated to any other condition are referred to as chronic or post-COVID-19 syndrome. A meta-analysis revealed that at least 45% of survivor of COVID-19 continued to have at least one residual symptom for an average of 126 days. The most common symptom recorded (28.4%–34.8%) was fatigue, which was followed by aberrant CT/X-ray patterns (45.3%), ground glass opacification (41.1%), and reduced carbon monoxide diffusion capacity (31.7%) (O'Mahoney et al., 2023).

Post COVID-19 syndrome patients may experience organ and tissue dysfunction as a result of systemic inflammatory response syndrome (SIRS). It is believed that PCS and Mast Cell Activation Syndrome are connected (MCAS). Stressor-induced interactions can lead to aberrant control of the activation of mast cell in PCS by inducing activation of mast cell genes via SARS-CoV-2 infection (Maxwell et al., 2021). Autoantibodies are produced during SARS-CoV-2 infection as a result of TLR activation on immune cells. Autoantibodies can also activate mast cells by interacting with mast cells' immunoglobulin receptors (Mukherjee et al., 2021). Following ACE2 expression, mast cells can release vasoconstrictor leukotrienes (LTs), which influence the pulmonary Renin-Angiotensin system (RAS). Several inflammatory mediators, such as histamine, prostaglandins (PGs), heparin, tryptase, Leukotrienes, and chemokines like Interleukin (IL)-1 β and IL-6, are released when mast cells are stimulated. Adrenocorticotrophic hormone (ACTH), cortisol, and corticotrophic releasing factor (CRF) are all released in greater amounts under stressful circumstances when infected with SARS-CoV-2. Moreover, intestinal stress alters the gut microbiota, which has an impact on tryptophan metabolism, neurotransmitter release, and short-chain free fatty acids (SCFAs) release. All of these changes lead to a stronger activation of mast cells, which in turn produces more inflammatory cytokines. Thus, mast cells provide neurological, immunological, and endocrine pathways that link the gut and the brain (Fuchs et al., 2012).

To preserve immunological homeostasis, a balanced anti-inflammatory response would typically be triggered by COVID-19's strong pro-inflammatory and overactive immune responses (Range and Moser, 2013; Sugimoto et al., 2016). This would result in an immunosuppressive condition. Nonetheless, chronic immunosuppression may result in PCS (Cañas, 2020). Transformation-promoting factor β (TGF- β) is an immunosuppressive, profibrotic, and anti-inflammatory cytokine that is raised during and after SARS-CoV-2 infection to reduce excessive pro-inflammatory responses. However, in PCS this regulation does not work

properly. Thus, there is ongoing inflammation. According to a number of research (Goërtz et al., 2020), infection persistence also causes long-term immunological activation. Multisystem Inflammatory Syndrome (MIS) is a result of persistent infection activating autoreactive T lymphocytes via antigen presentation by antigen-presenting cells. While pro-inflammatory cytokine levels and MIS may be correlated, the severity of the first SARS-CoV-2 infection is not. One possible explanation for the delayed start of MIS following SARS-CoV-2 infection is adaptive immune response dysregulation. Similarly, in 52% of PCS patients, B cell activation and antiphospholipid autoantibody synthesis were noted (Koné-Paut and Cimaz, 2020). When combined, immunological dysregulation, autoimmune mimicry, pathogen reactivation, persistence of inflammatory responses, and modifications to the host microbiome may lead to PCS. **Table 1** describes the pathophysiology of PCS in various organ systems.

4. Clinical Manifestation of Post COVID-19 Syndrome

Acute infection intensity and duration influence the clinical presentation of PCS, which can take many different forms. Fatigue is the most common PCS symptom, affecting 17–72% of critically COVID-19 patients. Respiratory symptoms such as dyspnea (10–40%), exercise intolerance (10–40%), and chest pain (22%), are more common in patients with PCS. Cardiovascular symptoms, including hypertension, arrhythmias, and postural hypotension, are brought on by endothelial dysfunction and heart injury (Mitrani, Dabas, and Goldberger, 2020). Thirty percent of COVID-19 patients may experience gastrointestinal symptoms for more than two months following hospital discharge, including nausea, vomiting, diarrhea, bowel movements, and appetite abnormalities (Carvalho-schneider et al., 2020). Neuropsychiatric problems, including smell/gustatory impairment, affect 9–11% of individuals and can last for six to eight months with mild COVID-19 (Sugiyama et al., 2022). Anxiety and depression account for 26% and 40%, respectively, of psychiatric consequences in PCS patients that manifest six months after infection (Alemanno et al., 2021). Aggression, cognitive impairments, decreased social engagement, obsessive-compulsive disorders, and stress disorders account for 43% of these difficulties. Guillain-Barre syndrome, stroke ischemic, seizures, transverse myelitis, and, cerebral vasculitis, are among the neurological conditions that have been seen in PCS patients (Scoppettuolo et al., 2020). **Table 1** and **Figure 1** describe the clinical manifestation of PCS in various organ systems.

In severe conditions, patients with PCS can develop some complications. Pulmonary complications, including lung fibrosis and radiological abnormalities that may persist for up to six months, are the most common complications among individuals with PCS. Furthermore, pulmonary dysfunction in PCS patients results in reduced gas exchange even with normal lung computed tomography (CT) scans. Due to lung fibrosis or scarring, patients with PCS may have dyspnea and a chronic cough (Batiha et al., 2022). Neurological problems encompassing musculoskeletal disorders and peripheral and central neurological disorders, accounted for around 36.4% of the patient. Concerning COVID-19, headache (13.1%), fatigue (14.0%), and dizziness (16.8%) were the most common neurological symptoms. Moreover, COVID-19 has been connected to ataxia, disorientation, seizures, stroke, and other central nervous system symptoms and consequences. Due to the extreme burden of a significant inflammatory response, PCS patients may also experience anxiety, depression, and psychosis.

Acute SARS-CoV-2 infection-related brain stem injury also causes cardio-respiratory dysfunction because it damages the respiratory and vasomotor areas Sinus tachycardia, postural orthostatic tachycardia syndrome (POTS), endothelial dysfunction, autonomic dysfunction, and dysautonomia can all occur in PCS patients. In addition, 5% of hospitalized COVID-19 patients get Acute Kidney Injury (AKI), and up to 31% of patients experience renal issues as a result of endothelial injury, inflammation, and coagulopathy. Gastrointestinal issues that patients may experience as a result of persistent GIT inflammation include reduced appetite, nausea, vomiting, diarrhea, and abdominal pain. Changes in the gut microbiota after infection may increase the risk of

lung dysfunction and systemic inflammation (Al-Buhadily et al., 2021)

Endocrine complications include thyrotoxicosis, thyroiditis, and Gravis' disease also can happen. In male COVID-19 patients, disturbance of the pituitary-testicular axis may result in decreased sperm production, quality, and testicular function. A case-control study found that 9.3% of participants had a low cortisol response and 46.5% had an insufficient growth hormone response. Thyroid stimulating hormone and prolactin levels also rose noticeably, by 9.3% and 4.6%, respectively. Some individuals also have increased peripheral insulin resistance in addition to decreased pancreatic cell insulin production. One of the hematological consequences that increases the risk of heart thrombus and ischemic stroke is thromboembolism (Moreno-Perez et al., 2022) .




Fig. 1. Pathophysiology of Post-COVID-19 Syndrome. This syndrome occurs in some patients who experience acute COVID-19 infection (1). The chronic inflammation that occurs in PCS is caused by the persistence of viral antigens and an autoreactive immune response as well as intestinal dysbiosis and microvascular dysfunction due to acute infection (2). As a result, multi-organ failure occurs (3).

Table 1. Pathophysiology of Post COVID-19 Syndrome and Its Complication in Organ Systems

Organ System	Pathophysiology(s)	Clinical Manifestation(s)	Reference(s)
Neurosensory	- Neuroinflammation is a chronic inflammatory process that activates brain microglia as a result of viral invasion or after the acute stage of the illness. - Coagulation activation: Microthrombosis hinders neuronal transmission and tissue vascularization. - Autoimmunity: atypical immune reaction during the acute stage of disease or molecular emulation of SARS-CoV-2 and host antigens. - Abnormalities in brain metabolism. - Remaining virus particles: support the low-level inflammatory reaction. - Nerve root and peripheral trigeminal nerve activity.	- Chronic headaches - Sleep disorders - Olfactory and gustatory impairments - Brain fog, memory and concentration problems - Depression, anxiety, post-traumatic stress disorder - Vertigo, imbalance, psychosis, and hallucinations - Small fiber neuropathy - Tremor in the posture - Pain syndromes - Neurodegenerative disorders - Poor concentration	(Al-Buhadily et al., 2021; Castanares-Zapatero et al., 2022; Mehandru and Merad, 2022)
Musculoskeletal	- External variables, such as musculoskeletal conditions. - Environmental factors: humidity, temperature, and social isolation. - concomitant conditions. - Congestion in the lymphatic-lymphatic system. - Muscle bioenergetic diseases resulting from dysfunctional mitochondria.	- Hearing loss - Tinnitus - Red eyes - Sore throat - Fatigue - Myalgia - Joint pain - Small joint arthritis	(Mehandru and Merad, 2022) (Al-Buhadily et al., 2021; Castanares-Zapatero et al., 2022; Mehandru and Merad, 2022)
Immune system	- Myocytes disruption and activation of fibroblast. - Alterations in the bone's microcirculation, including leukocyte aggregation, hypercoagulability, and vascular inflammation. - Tension in joints and activated immunity. - Chronic disorganized immune system activation: low-grade inflammation leading to multiple organ dysfunction.	- Altered immune system - Increased risk of infection	(Mehandru and Merad, 2022)

Gastrointestinal and hepato-biliary system	• Mast cell activation syndrome • Persistent infection • Multisystem inflammatory syndrome in children (MIS-C).		
	• Intestinal dysfunction following an infection • Alterations in the microbiota; and harm to the liver and biliary system. • Autonomic nervous system abnormalities (intestinal motility disorders).	• Appetite loss • Pain in the abdomen • Nausea • Loss of weight • Changes in bowel motility • Irritable bowel syndrome • Dysphagia	(Mehandru and Merad, 2022)
Renal system	• Invasion of renal cell virus • The severity of prior severe illness • Microangiopathy • The renin-angiotensin-aldosterone pathway is disrupted • The Glomerularopathy.	• Microhematuria	(Mehandru and Merad, 2022)
Endocrine system	• Pancreatic β cell invasion by a virus • Direct injury to the thyroid gland, subacute thyroiditis, low T3 syndrome.	• Subacute thyrotoxicosis • Persistent glycemic abnormalities • Grave's diseases • Hashimoto's thyroiditis • Lipid abnormalities	(Mehandru and Merad, 2022)
Cardiovascular	• Myocardial injury can occur directly from viral infection, which is thought to be less prevalent, or indirectly through systemic inflammatory response. • Desmosomal protein fibro-fatty displacement and cardiomyocyte death. • Corticosteroids and renin-angiotensin-aldosterone system (RAAS) imbalance.	• Chronic headaches • Sleep disorders • Olfactory and gustatory impairments • Brain fog, memory and concentration problems • Depression, anxiety, post-traumatic stress disorder • Vertigo, imbalance, psychosis, and hallucinations • Small fiber neuropathy • Tremor in the posture • Pain syndromes • T2 signal and positive LGE • Myocardial oedema • Pericardial effusion	(Visco et al., 2022)

Respiration system	• An abnormal and persistent inflammatory reaction • An excessive fibroblastic growth • Leukocyte telomere length and lactate dehydrogenase levels as separate risk factors for the emergence of fibrotic-like abnormalities	• Hemosiderin deposits • Diastolic dysfunction • Capillary abnormalities • Arrhythmias • Pulmonary hypertension • Dyspnea • Persistent cough • Asthma exacerbation • Reduced carbon monoxide (DLCO) diffusing capacity • Dyspnea, tachypnea, pleurisy; cough • Disturbances in lung function ($\downarrow$ FEV1, $\downarrow$ FEV1/FVC) • Pulmonary fibrosis • Crazy paving in the interstitial space • Ground-glass opacity residual • Abnormal diffusion • Pulmonary embolism • Pneumonia	(Visco et al., 2022)
Dermatological	• Immune responses • Epitope spreading. • Cryptic antigens are presented to functional T cells. • The phenomenon of antibody-dependent enhancement (ADE) is observed when human hair follicle dermal papilla cells are directly attacked by viruses. • The production of microthrombi in the coagulation cascade might block the blood supply to hair follicles. • Underlying pro-inflammatory cytokines that are released as a result of chronic inflammation. • Stress in the acute stage (stress related to emotions, fever).	• Skin rash • Telogen effluvium • Nail alterations	(Mehandru and Merad, 2022; Monari et al., 2022)

Systemic manifestation	• The organs' modified homeostatic environment (stromal damage, less tissue-resident macrophages). • Proinflammatory cells that persist. • Modified generation of cytokines. • Modified immunometabolic routes (heme-related metabolic pathways, ROS). • The altered gut bacteria, epitope dissemination, and immunological antibodies.	• Chronic malaise • Restriction of daily activities • Asthenia	(Mehandru and Merad, 2022)
Laboratory abnormality	• Long-lasting proinflammatory cells that infiltrate neutrophils, plasma cells, effector T cells, monocytes, and PDCs • Modified production of cytokines (IFN-α Type I, IFN-γ, and III, IL-6, TGF-B, and IL-1B) • Modified immunometabolic routes (heme-related metabolic pathways, ROS) • autoimmune Antibodies • The spreading epitope • Changes in intestinal microbiota	• Increased neutrophils, hepatocyte growth factor (HGF), matrix metalloprotease 7 (MMP-7), and lipocalin 2 (LCN-2) • Increased D-dimer levels and anemia • Anomali of lipids • Abnormal liver function tests • Thrombocytosis • Elevated hemoglobin A1C (HbA1C) • Reduced serum albumin • Coagulation abnormalities • Electrolyte abnormalities	

5. Recent Alternatif Therapy for Post COVID-19 Syndrome

5.1. Herbal Medicine

One of the most popular uses of herbal medicines is in Traditional Chinese Medicine (TCM). Traditional Chinese medicine (TCM) has long been used to treat a wide range of ailments in both home and foreign countries. For instance, it has been shown that Lianhuaqingwen (LH), a Chinese medicine made up of thirteen herbs, efficiently cures a variety of influenza viruses by regulating the immune system, preventing the spread of the virus, and reducing the length of time an H1N1 infection lasts. As a therapy, it is just as effective as oseltamivir (Xiong et al., 2020). Moreover, a retrospective analysis of from patients who acquired SARS-CoV-2 at Wuhan Ninth Hospital and General Hospital has shown that Lianhuaqingwen is helpful in the treatment of the virus. Combining LH can greatly lessen the severity of COVID-19 illness and ameliorate cardinal symptoms (Runfeng et al., 2020).

India is among the nations with the most advanced systems of herbal medicine, aside from China. For many years, Ayurveda has been the main medicine utilized to control many ailments. The antiviral and immunosuppressive properties of Ayurveda on T and B lymphocytes are known to lessen the inflammatory effects of COVID-19 infection (Dhane and Shinde, 2022). The M protein of SARS CoV-2 and other target proteins are susceptible to the antiviral effects of certain Ayurvedic ingredients, such as *Tinospora cordifolia*, *Withania somnifera*, and *Ocimum sanctum*, according to molecular docking research. Ayurvedic phytoconstituents (Maurya and Sharma, 2022) may block the SARS Corona Virus's papain-like protease and primary protease, as well as cause disruptions to the relationship between the host receptor ACE 2 and the viral S-protein RBD (Receptor Binding Domain) (Balkrishna et al., 2020; Shree et al., 2022).

In addition to providing relief during the acute stage of COVID-19 infection, a number of herbal remedies have demonstrated beneficial effects in PCS, particularly in easing lingering symptoms following negative COVID-19 test results. A number of active ingredients in herbal remedies, including green tea, kadha, and tulsinol from *Ocimum sanctum* (tulsi), actively work to minimize the onset of symptoms by inhibiting the NF- κ B pathway and pro-inflammatory cytokine production (Kumar et al., 2020). The constituents, mechanisms of action, and therapeutic effects of various herbal medicines in treating PCS symptoms are presented in **Table 2**.

Compared to chemical treatments, herbal remedies seen more "natural" and "safer". However, as the body of research on herbal remedies grows, a growing number of side effects have been documented. These include reactions to the biologically active components of herbs, side effects brought on by contaminants, and chemical interactions between herbs and drugs. Gastrointestinal disturbances, including aberrant liver function, renal dysfunction, headache, nausea, vomiting, diarrhea, and appetite loss, are the most frequent side effects of herbal therapy (Wang et al., 2022). One of the conditions that is frequently linked to the usage of some Chinese herbs, such as *Aristolochia fangchi*, is nephropathy. The presence of pyrrolizidine alkaloids, which are unintentionally or intentionally added to herbal remedies, is another frequent cause of toxicity for these remedies. This alkaloid induces hepatotoxicity and eventually results in deadly veno-occlusion.

Table 2. Herbs Component and Mechanism of Action of Several Patent Herbal Medicine

Herbal Medicine (s)	Origin	Theraupetic Effect(s)	Herb(s) component	Mechanism of Action(s)	Reference(s)
Jinhua Qinggan (JHQG) granules	Traditional Chinese Medicine (TCM)	- JHQG proved highly successful in combating the avian influenza A (H1N1) virus, for which it was advised. - JHQG may also help influenza patients' immune systems work better. - Empirical research indicates a positive impact on symptoms related to exhaustion, persistent cough, diarrhea, heightened appetite, chest discomfort, and muscle soreness.	<i>Gypsum Fibrosum</i> , <i>Lonicera japonica</i> , <i>Prunus sibirica</i> L., <i>Ephedra sinica</i> , <i>Scutellaria baicalensis</i> , <i>Fritillaria thunbergii</i> , <i>Forsythia suspensa</i> , <i>Anemarrhena asphodeloides</i> , <i>Artemisia annua</i> L., <i>Arctium lappa</i> L., <i>Mentha haplocalyx</i> Briq., and <i>Glycyrrhiza glabra</i> L.	Anti-Inflammations - Prevents the release of cytokines that promote inflammation (IL-6, TNF-α, IL-7, IL-1β, IL-4, IL-2 and IL-13). - Prevents the release of macrophage chemokine protein-1, IL-6, TNF-α, and promotes protein-10. - Inflammatory signaling pathways: the PI3K/Akt signaling pathway, Janus kinase/signal transducer and activator of transcription (JAK/STAT) signaling system, MAPK pathway, and nuclear factor kappa B (NF-κB), are blocked.	(Xia et al., 2020; Cao et al., 2022; Zhao et al., 2021; Wang et al., 2022; Zhu et al., 2023; Zhao J et al., 2021; Xiong et al., 2022; Cruz et al., 2021; Xu et al., 2021; Li et al., 2022; Panda et al., 2021; Yates et al., 2022; Dhane and Shinde, 2022; Rastogi et al., 2022; Anderson et al., 2019)
Lianhua Qingwen (LHQW) capsules	TCM	- LHQW is advised in addition to Western medication for the treatment of influenza, particularly avian influenza A (H1N1). - RCT studies demonstrate an improvement in the symptoms of muscle soreness, increased hunger, persistent cough, and diarrhoea.	<i>Ephedrae Herba</i> , <i>Armeniacae Semen Amarum</i> , <i>Lonicerae Japonicae Flos</i> , <i>Glycyrrhizae</i> , <i>Forsythiae Fructus</i> , and <i>Menthae Haplocalcis Herba</i> , <i>Atractylodis Rhizoma</i> , <i>Pogostemonis Herba</i> , and <i>Gypsum Fibrosum</i>	- Inhibits the TLR4 – Myd88 – NFκB pathway. - Decreases the generation of monocytes, neutrophils, lymphocytes, and white blood cells. - Reduces prostaglandin E2 (PGE2) synthesis in the vicinity of the hypothalamus. - Restricts cyclooxygenase-2 (COX-2) expression.	
Huashibaidu (HSBD) granules	TCM	- The treatment of epidemic toxin blocking in the lung, a pattern observed in severe COVID-19 cases, has been suggested to use HSBD. - Greatly reduces weariness, cough, and chest discomfort. eases dry mouth and headaches.	<i>Glycyrrhizae Radix et Rhizoma</i> , <i>Gypsum Fibrosum</i> , <i>Ephedrae Herba</i> , <i>Ageratum rugose</i> , <i>Atractylodis Rhizoma</i> and <i>Descurainiae Semen Lepidii Semen</i> , <i>Poria</i> , <i>Pogostemonis Herba</i> , <i>Armeniacae Semen Amarum</i> , <i>Tsaoko Fructus</i> , <i>Pinelliae Rhizoma</i> <i>Praeparatum</i> , <i>Astragali radix</i> , and <i>Paeoniae Radix Rubra</i>	- Represses the expression of TLR-4 produced by LPS and reduces the inflammatory cellular pathway cascade mediated by TLR. - Prevents the synthesis of NO, iNOS, COX-2, IL-6, and TNF-α. - By preventing the synthesis of TXB2 and LTB4 metabolites, it lowers the control of COX, 5-LO, and 5-HETE.	
Xuanfeibaidu (XFBD) granules	TCM	- The primary goal of treating COVID-19 individuals who are moderately to severely ill. - Reduction of fever, shortness of breath, cough, and exhaustion.	<i>Ephedrae Herba</i> , <i>Glycyrrhizae Radix et Rhizoma</i> , <i>Armeniacae Semen Amarum</i> , <i>Atractylodis Rhizoma</i> , <i>Descurainiae Semen Lepidii Semen</i> , <i>Atractylodis Rhizoma</i> , <i>Pogostemonis Herba</i> , and <i>Gypsum Fibrosum</i> .	Blocks the Wnt/β-catenin and ERK pathways.	

Lung Cleansing and Detoxifying Decoction (LCDD) (consists of Ma Xing Shi Gan (MXSG); She Gan Ma Huang decoction (SGMH), Xiao Chai Hu (XCH), and Wu Ling San (WLS))	TCM	• The management of H1N1 influenza. • The management of tonsillitis, asthma, and flu-like symptoms. Treatment of COVID-19 individuals, both severe and non-severe, is advised. • Relieving early-stage extrapulmonary symptoms of diarrhea or problems related to the renal, hepatic, or cardiac systems. decreased sore throat, cough, and weariness.	<i>Asari Radix Et Rhizoma, Belamcandae Rhizoma, Dioscoreae Rhizoma, Farfarae Flos, Aurantii Fructus Immaturus, Asteris Radix et Rhizoma, Armeniacae Semen Amarum, Citri Reticulatae Pericarpium, Zingiberis Rhizoma Recens, Scutellariae Radix, Cinnamomi Ramulus, Ephedrae Herba, Pinelliae Rhizoma, Tuckahoe, Herba Agastachis, and Glycyrrhizae Praeparata cum Melle Radix et Rhizoma</i>	Antioxidants • Boosts the activity of superoxide dismutase, Ca2plus-ATPase, and Naplus-Kplus-ATPase, therefore lowering free radicals; • Decreases oxidative stress indicators (H2O2 and NO). • Effectively combats hydroxyl and DPPH free radicals • Anti-lipoxygenase and anti-proteinase properties. • Reverse NADPH oxidase overexpression 4. The ERK/Nrf2/HO-1 signaling pathway is upregulated. • Preventing intracellular ROS from building up. strong DPPH-lowering action. • Higher glutathione and lower amounts of malondialdehyde in the lungs.
Banlangen granule (BLG)	TCM	• Detoxifying and therapeutic agent for exogenous fever, carbuncle sores, erysipelas, mumps, and epidemic heat syndrome. • Treatment, management, and prevention of respiratory viral infections (viral hepatitis, influenza A subtype H1N1, and viral skin disorders). • The method most frequently employed to stop COVID-19 infections.	<i>Radix Isatidis</i>	Immunomodulators • TNF, IFN-γ, C-reactive protein (CRP), peroxisome proliferator-activated receptor-gamma, and tumor protein P53 are examples of pro-inflammatory cytokines that are modulated. • Modulates T helper cell differentiation (Th 1, Th 2, and Th 17 and T lymphocyte receptor pathways. • Enhances thymus and spleen function. • Promotes the growth of CD4+ and CD8+ T cells. • Increases the expression of IgM and IgG. • Glycyrrhizin stimulates macrophages and natural killer (NK) cells via the signaling pathways of Toll-like receptors (TLRP) and Mitogen-Activated Protein Kinase (MAPK).
Qu Du Qiang Fei I Hao Fang (QDQF1)	TCM	• Treat and prevent infections in the upper respiratory tract caused by influenza H1N1. • Address bleeding, chronic weariness, anemia, appetite loss, phlegmatic problems, and weakness.	<i>Radix Glycyrrhizae, Atractylodis Macrocephalae, Lonicerae Japonicae, Fructus forsythi, Astragali Radix, Saposhnikoviae Radix, Lonicerae Japonicae, and Forsythiae Fructus)</i>	
Shufeng Jiedu	TCM	• Treatment for acute lung injury	<i>Bupleurum chinense, Fallopia japonica, Forsythia</i>	

capsule (SFJD)		(ALI) and respiratory infections. - Treatment influenza A (H1N1)-related upper respiratory tract - Treatment for Common cold, Acute Rhinitis, Herpangina, Acute pharyngitis, and Acute tonsillitis.	<i>suspensa</i> , <i>Glycyrrhiza uralensis</i> , <i>Isatis indigotica</i> , <i>Patrinia scabiosaefolia</i> , <i>Phragmites australis</i> , and <i>Verbena officinalis</i>	- Macrophages' phagocytic activity towards andrographolide - Boost host cells' cellular immune response. - Thoothuvalaini increases antioxidant levels and inhibits lipid peroxidation enzymes in the blood and liver.
Jing Fang Bai Du pill (JFBD)	TCM	- Management of common cold, exogenous febrile illness, widespread epidemic infections, headache, lightheadedness, restlessness, body aches, cough from lung cold, and other symptoms. - Epidemic infections, common cold, diarrhea, sores, and influenza are among the indications for cold dampness.	<i>Bupleurum chinense</i> DC, <i>Ligusticum chuanxiong</i> Hort., <i>Angelica pubescens</i> , <i>Saposhnikovia divaricata</i> , <i>Poria cocos</i> , <i>Glycyrrhiza uralensis</i> , <i>Schizonepeta tenuifolia</i> , <i>Platycodon grandiflorum</i> (Jacq.), <i>Peucedanum praeruptorum</i> , <i>Notopterygium inchum</i> , and <i>Citrus aurantium</i> L.	- In host cells, flavonoids promote the production of interferons that fight viruses. - Inhibition of B-lymphocyte proliferation, elevation of hemagglutination titers, and reduction of serum IgA, IgM, and C3 levels. - Focus on controlling the PD-1/IL17A axis to control the invasion of macrophages and neutrophils. - Inhibition of the potentiation of delayed type hypersensitivity (DTH) reaction brought on by cyclophosphamide.
Qingfei Paidu decoction (QFPD)	TCM	- Suggested for the management of mild to severe COVID-19 - Hasten the resolution of cough symptoms; reduce exhaustion; and stimulate appetite.	<i>Ephedra sinica</i> , <i>Glycyrrhiza uralensis</i> , <i>Semen Armeniacae</i> , <i>Gypsum Fibrosum</i> , <i>Cinnamomum cassia</i> , <i>Alisma plantago-aquatica</i> ., <i>Polyporus umbellaru</i> , <i>Atractylodes Macrocephala</i> , <i>Poria cocos</i> , <i>Bupleurum chinensis</i> DC., <i>Scutellaria Baicalensis</i> Georgi., <i>Pinellia ternata</i> Zingiber officinale, and <i>Aster tataricus</i>	Enhanced PMN cells, including monocytes and macrophages, as well as neutrophils, eosinophils, mast cells, and basophils.
Ayush-64	India Alternative and Complementary Therapies (IACT)	- Reduced cough, dyspnea, pyrexia, headache, and sore throat. - Commonly used for the common cold, sore throat, cough, runny nose, fever, and other illnesses.	<i>Alstonia scholaris</i> R. Br., <i>Picrorhiza kurroa</i> Royle ex. Benth, <i>Swertia chirata</i> Pexbex. Karst, and <i>Caesalpinia crista</i> L., <i>Ocimum sanctum</i> , <i>Cinnamomum verum</i> , <i>Zingiber officinalae</i> , <i>Piper nigrum</i> .	Organ Protections - Decreased apoptosis caused by LPS thus reducing ALI incidence. - It reduces pulmonary oedema, inhibited inflammatory cell infiltration and the over-expression of inflammatory cytokine. - Inhibit fibroblast collagen Improve intestinal flora. - Decrease cough sensitivity, iron chelating agent, scavenges hydroxyl radicals, superoxide radicals, and DPPH radicals. - Relieve ox-LDL-induced oxidative stress via the ERK/Nrf2/HO-1 signalling
Ashwagandha	IACT	Well known as a rejuvenator drug indicated in dyspnea and cough.	<i>Withania somnifera</i> (L.) Dunal	
Guduchi	IACT	Hepatoprotective, antimicrobial,	<i>Tinospora cordifolia</i> Willd.	

Ghanavati		antioxidant, antitoxic, antidiabetic, hypolipidemic, anti-malarial, and anti-neoplastic.	
Chyawanprasha	IACT	It is used to treat conditions like asthma, TB, allergic bronchitis, the common cold, and cough.	<i>Phyllanthus emblica</i>
Nilavembu Kudineer (NVK) decotion	IACT	Antiviral for Chikungunya and Dengue virus.	<i>Andrographis paniculata</i> , <i>Vetiveria zizanioides</i> L., <i>Santalum album</i> L., <i>Piper nigrum</i> L., <i>Cyperus rotundus</i> L., <i>Hedyotis corymbosa</i> L.Lam., <i>Plectranthus vettiveroides</i> (K.C.Jacob), <i>Zingiber officinale</i> Roscoe., and <i>Trichochanthes cucumerina</i> L.
Kaba Sura Kudineer (KSK) decotion	IACT	- It is commonly used for fever and respiratory diseases. It's used to treat pox viral infections and herpes infections. - Accelerate the disappearance of cough, shortness of breath, breathlessness, and fever.	<i>Piper longum</i> L., <i>Anacyclus pyrethrum</i> L., <i>Zingiber officinale</i> Roscoe., <i>Tragia involucrate</i> L., <i>Solanumanguivi</i> Lam., <i>Anisochilus carnosus</i> , <i>Costus speciosus</i> , <i>Tinospora cordifolia</i> , <i>Syzygium aromaticum</i> , <i>Clerodendrum serratum</i> (L.), <i>Andrographis paniculata</i> (Burm.f.), <i>Cyperus rotundus</i> L., <i>Terminalia chebula</i> , <i>Justicia adathoda</i> L., and <i>Sida acuta</i> .

pathway, NOS, NO, and MDA activity.

- Promote the growth and regulation of normal flora in the gut, such as *Bifidobacteria*, *Lactobacilli*, and *Lactococcus*.
- Suppress alveolar epithelial apoptosis initiated via NF-κB and p38 MAPK pathway.
- Decreases total neutrophil counts and helps clear the lungs, preventing lung impairment—inhibition of Angiotensin-Converting Enzyme II (ACE2).
- Activating a transcription factor (Nrf)–2 causes evading apoptosis in an Nrf-2-dependent manner in HL-1 cardiomyocytes.
- Reversed the effects of ischemia-reperfusion on the activities of lipid peroxidation (LPO), catalase (CAT), glutathione peroxidase (GPX), and superoxide dismutase (SOD).

5.2. Traditional Practices

- **Acupuncture**

For many years, several problems have been addressed using acupuncture. The medical philosophy and treatment technique of acupuncture is predicated on the idea of applying pressure or tiny needles to particular body locations. Acupuncture aims to balance surplus energy that is flowing through specific body channels (Song et al., 2019). Acupuncture is often recommended for the treatment of pain, both acute and chronic, including fibromyalgia, rheumatoid arthritis, chronic daily idiopathic pain, migraine, tension headaches, myofascial pain syndrome, and nausea and vomiting (Schweiger et al., 2020). It is believed that acupuncture reduces inflammation (Kwon et al., 2018). Nonetheless, two particular contraindications deserve special attention: those who are expected to have psychosis or delusions, as well as patients who have an automatic implanted cardioverter-defibrillator (AICD) (Zhang J et al., 2022).

Additionally, acupuncture can lessen the intensity of anxiety and sadness, enhance sleep, treat insomnia, and improve cognitive damage following a stroke. Acupuncture also addresses physiology, neuropsychiatric problems, and olfactory impairment (anosmia, ageusia, and sleeplessness) (Liu et al., 2020). Acupuncture has the potential to be beneficial in treating some contemporary acute infectious diseases, acute bacterial dysentery, epidemic dengue fever, and severe acute respiratory syndrome (SARS) (Xu et al., 2021; Wang et al., 2020). Its benefits include fever reduction, phlegm and cough relief, gastrointestinal symptom relief, respiratory function improvement, and prevention of disease progression (Helianthi et al., 2022).

In mild-to-moderate COVID-19 patients, the combination of acupuncture and pharmaceutical medications has already shown promise, as evidenced by a noteworthy reduction in the length of cough symptoms compared to solo pharmaceutical therapy (Boehmer et al., 2020). Acetylcholine is released more when vagus cholinergic anti-inflammatory pathway is activated by acupoint stimulation. After binding to the $\alpha 7nAChR$ receptor on macrophages, acetylcholine prevents the release of cytokines that promote inflammation. The anti-inflammatory adrenal-dopamine vagus-medullary pathway is triggered by stimulation of other sites, resulting in the production of dopamine. This neurotransmitter can bind to type I dopamine receptors and suppress pro-inflammatory cytokines such as IL-6, INF- γ , and TNF α . When acupoints are stimulated simultaneously, TLR-4 and NF- κB activation is inhibited, which lowers the release of pro-inflammatory cytokines (Boehmer et al., 2020; Wang and Pariente, 2021). In addition, it has been demonstrated that stimulation at sites LI4 and ST36 raises blood levels of cortisol and ACTH and directly suppresses the production of IL-6, which lowers ferritin and CRP. Additionally, acupuncture can enhance host-disease interactions and lessen the generation of ROS and NO (Morita et al., 2022).

According to clinical studies, patients were treated by two TCM-licensed acupuncturists (LAc) who inserted auricular needles with insertion depths ranging from 0.1 to 1.0 cun before receiving body needles. Thirty minutes retention of needle in conjunction with De Qi, twisting, and lifting techniques has been shown to improve various PCS symptoms, including pain, ageusia, fatigue, anxiety, anosmia, brain fog, dyspnea during exercise, dry cough, chest pressure, and palpitations (Trager et al., 2022; Hampton and Bartz, 2021). No adverse effects have been reported.

- **Yoga**

Yoga is a comprehensive kind of physical and mental well-being that has its roots in ancient India. Numerous yoga practices, including asana (postures), dhyana (meditation), and pranayama (breathing exercises), seem to provide a variety of therapeutic benefits. Yoga has been shown to improve mental health, including lowering anxiety and sadness, as well as metabolic problems and low back pain (Saeed et al., 2019). Energy balance is achieved by yoga practice, and immune system performance depends on it. The hypothalamus's posterior or sympathetic region may be inhibited by yoga. This inhibition

optimizes the body's sympathetic response to stressful stimuli and restores the autonomic regulating reflex processes associated with stress. While blocking the regions in charge of anger, violence, and fear, yoga practice activates the positive pleasure centers in the middle forebrain and other areas that contribute to moods of contentment and pleasure. Reduced anxiety, heart rate, breathing rate, blood pressure, and cardiac output are the results of this inhibition. Regular yoga practice lowers levels of monoamine oxidase, an enzyme that degrades cortisol and neurotransmitters, and dramatically raises serotonin levels. This helps people get over their worry, tension, and despair (Bushell et al., 2020). Another advantage of yoga is increased mobility, which is achieved by increasing relaxation of the muscles and connective tissue around the bones and joints with continued practice. This is thought to be one of the reasons yoga is associated with reduced aches and pains. As a result, yoga has been suggested as a kind of rehabilitation for individuals who have had COVID-19 or who have symptoms of PCS. Through its anti-inflammatory, anti-oxidant, and immunomodulatory qualities as well as its ability to increase serotonin activity, yoga exerts organ-protective benefits on the cardiovascular and respiratory systems (Basu-Ray et al., 2022; Srinivasan et al., 2021).

In a clinical trial, quality of life, pulmonary ventilation, and muscle strength—particularly biceps and quadriceps—can all be enhanced by pursed lip breathing combined with bhasrika pranayama exercise three times a day for five minutes each session for six weeks. No negative effects have been documented (Macedo et al., 2023). An additional workout routine consists of 60 minutes of yoga practice twice a week for eight weeks. The breathing exercises consist of three parts: ten minutes of standing breathing, ten minutes of sitting breathing with an asana, and ten minutes of stretching breathing. These exercises have been shown to enhance sleep quality, lessen anxiety, depression, and stress episodes, relieve dyspnea, improve lung function and capacity, increase vital capacity, lower heart rate, lower systolic blood pressure, and raise tidal volume.

- **Tai Chi**

Chinese martial art tai chi is a form of light to moderate cardiovascular exercise that integrates mental and physical training. Tai Chi is one among the suggested physical exercises during the COVID-19 pandemic to enhance fitness, including boosting heart rate, blood pressure, body mass index, weariness, mental health, sleep difficulties, balance and motor control, and strength and aerobic capacity (Easwaran et al., 2021; Polero et al., 2021; Shu et al., 2021).

Tai Chi is an indirect anti-inflammatory and immunomodulatory agent since it can alter the immune system and pro-inflammatory cytokine responses. Both the innate and adaptive immune systems' cell counts can be raised by practicing tai chi. Engaging in regular Tai Chi exercise raises T cells and the percentage of natural killer (NK) cells, which raises the Th1 immune response and may improve antiviral function. Furthermore, Tai Chi can decrease CRP as well as proinflammatory cytokines like IL-1b, TNF- α , and IL-6 (Zouwt et al., 2018). Tai chi also promotes the synthesis of other enzymes, including MnSOD, iNOS, and glutamylcysteine synthetase, which has an indirect antioxidant impact. By controlling elevated levels of cortisol, adrenaline, and insulin-like growth factor and boosting the synthesis of brain-derived neurotrophic factor and insulin-like growth factor—two neurobiological markers linked to emotional regulation—tai chi practice also lessens negative emotions like depression, anxiety, and melancholy (Zhang et al., 2019; Wu et al., 2022).

Five Tai Chi moves that were applied by the classic Cheng Man-Ch'ing Yang-style short form were used in several therapeutic investigations. For a duration of 12 weeks, patients participated in twice-weekly tai-chi courses, lasting roughly 60 minutes each. The outcomes of these studies demonstrate that PCS patients improve lung function, including FEV1/pred and FEV1/FVC, by strengthening their respiratory muscles and expanding their lung capacity. Diminish unfavorable feelings including despair, worry, and depression. improved bodily health statement, quality of sleep, and symptoms of exhaustion.

5.3 Nutraceutical and Supplementations

Anti-inflammatory, immunomodulatory, antiviral, and antioxidant compounds are known as nutraceuticals. Nutraceuticals frequently contain zinc, vitamins, curcumin, probiotics, selenium, and quercetin, among other ingredients, according to Barrea et al., 2022. These ingredients might provide alternative therapeutic and prophylactic effects for PCS and COVID-19 (Fratta et al., 2021; Pastor et al., 2021). Nutraceuticals are made up of active phytochemical components that are isolated from plants, food supplements, and functional foods. They are known to have medicinal properties (Catalano et al., 2022).

- **Vitamins**

One of the supplements that is advised for PCS prevention, treatment, and recovery is vitamin C. It is well-recognized that some vitamins, including A, C, D, and E, can treat PCS and COVID-19 disorders. It has been demonstrated that ascorbic acid, sometimes referred to as vitamin C, has antithrombotic, anticancer, and antioxidant qualities. In general, fruits like oranges and kiwis as well as vegetables like spinach contain vitamin C (Zhao et al., 2021). Vitamin C can prevent thrombotic effects, minimize oxidative stress, lower inflammation, manage cytokine storms, and lessen vascular and alveolar damage, according to several clinical investigations on COVID-19 infection. Additionally, individuals with severe and serious COVID-19 showed a significant rise in CD4, lymphocyte count, and C-reactive protein (CRP) after administering vitamin C (Zhang J et al., 2021). A placebo-controlled pilot experiment by Zhang et al. (2021) demonstrated a substantial reduction in mortality with high-dose intravenous vitamin C (HDIVC) treatment, using a daily dose of 24 g of ascorbate. HDIVC does not influence the length of ventilation discharge days in patients, although it may enhance lung function as indicated by an increase in the PaO₂/FiO₂ ratio (Tosaco et al., 2022). Giving 500 mg of liposomal vitamin C and l-arginine for 28 days improved endothelial function, muscle strength, gait performance, and fatigue in patients with PCS who had persistent fatigue, according to another RCT study. The patients also reported feeling less tired than before. Therefore, one may consider using this supplement to enhance athletic performance and lessen persistent COVID-19 symptoms.

Furthermore, vitamin A has a clinical effect on weakness, tiredness, fever, and bodily aches. Along with improvements in symptoms like chills, coughing, shortness of breath, hyposmia, anorexia, and diarrhea, ten days after the intervention there was also a drop in the amount of C-reactive protein as well as white blood cell as compared to normal treatment alone (Rohani et al., 2022; Masnadi et al., 2021b). Furthermore, vitamin A plays a crucial role in the formation of helper T cells, T cells, and B cells, all of which are essential for the maintenance of the immune system (Albergamo et al., 2023).

Clinical investigations and recent meta-analyses have shown that vitamin D functions by altering the expression of ACE 2 and the renin-angiotensin system (RAS). This will induce the ACE2/Ang axis, 1-7, and press the ACE/Ang II/AT1R axes, hence reducing acute lung injury caused by lipoprotein (LPS). Furthermore, vitamin D stimulates defense systems like macrophages and lowers proinflammatory cytokines while raising anti-inflammatory cytokine levels, which can eradicate SARS-CoV-2. It also stimulates the creation of natural antimicrobial peptides. A clinical trial (Fernandes et al., 2018) found improvements in ferritin, lactate dehydrogenase, IL-6, CRP, and the ratio of neutrophils to lymphocytes in patients who received 60,000 IU of vitamin D supplements daily for eight days. Although vitamin D3 had a positive effect on COVID-19 infection, the findings of Fernandes et al. 2018 and Popescu et al., 2022 did not justify the use of a single dosage of 200,000 IU for the repair of cytokines, chemokines, and growth factors. Nevertheless, given that a number of other studies have indicated that a vitamin D deficit is a predictor of a bad prognosis for COVID-19 patients, supplementing with adequate levels of vitamin D can still provide the therapeutic benefits of immunomodulators (Fratta et al., 2021). Furthermore, a comprehensive approach to treating Telogen effluvium, also known as post-infection hair loss, is necessary to address one of the side effects of PCS. This approach involves the use of zinc, biotin, vitamins A, C, E, and D, as well as organic compounds like Omega-3 (Magdalena et al., 2022).

There are high concentrations of vitamin E in green leafy vegetables, almonds, and sunflower and

olive oils. It's common knowledge that vitamin E is a strong antioxidant. As an antioxidant that protects cell membranes from oxidative stress, the component α -tocopherol has a defensive role. Because α -tocopherol may absorb reactive oxygen species (ROS), it can stop the oxidative burst that is linked to SARS-CoV-2. In addition, vitamin E promotes T-cell differentiation and proliferation, reduces the generation of inflammatory cytokines, and alters the inflammatory response in a number of organs, including the lungs. Studies with both human and animal models have shown that vitamin E enhances the immune response through nitric oxide production reduction, prostaglandin E2 reduction, cyclooxygenase-2 inhibition, T-lymphocyte communication, and modification of the Th1/Th2 ratio. Additionally, it controls protein kinase C to function as an immunoregulator.

- Mineral

Zinc (Zn) is the second most prevalent metal in the human body. Moreover, zinc has antiviral, anti-inflammatory, immune-stimulating, and membrane-stabilizing properties. In the bone marrow, zinc stimulates the development of B lymphocyte cells. In COVID-19, zinc inhibits RNA polymerase activity. Zinc responds well to low dosages of hydroxychloroquine, which also reduces the number of patients who need to be admitted to the hospital (Rohani et al., 2022). The effects of type-I interferon (IFN) can be amplified by zinc. Supplementing with zinc gluconate reduces the influx of neutrophils into the airways, releases TNF- α , and prevents the transcription of inflammatory genes that are dependent on NFkB. By reducing inflammatory cytokines (IL-6 and IL-1 β), zinc supplementation may help improve a protective type-I IFN response in COVID-19 (Pal et al., 2021).

- Fatty acids

Omega-3 polyunsaturated fatty acids with a long chain, also known as n-3 PUFA, have been shown to have anti-inflammatory, antioxidant, and psycho-neuro-immunological properties. Omega-3 polyunsaturated fatty acids have been demonstrated to treat mood disorders by decreasing proinflammatory cytokines, reestablishing the HPA axis, changing the gut-brain axis, modulating nerve transmission via lipid rafts, and maybe improving immunity to both physical and mental effects. Treatment with UFA omega-3 is advised for neuropsychiatric issues brought on by PCS, such as brain fog, weariness, anxiety, and depression. Strong anti-inflammatory properties are shown by the D-series resolvins, maresins, and protectins from DHA as well as the E-series resolvins from the Omega 3 group EPA. These properties inhibit the migration of neutrophils and monocytes across epithelial cells, as well as the production of cytokines (TNF- α and IL-1 β) and chemokines in wounded tissue. Furthermore, they facilitate the removal of leukocytes, apoptotic cells, polymorphonuclear neutrophils (PMN), and cell remains from inflammatory areas. Omega-3 polyunsaturated fatty acids (PUFAs) can directly reduce free radical levels because they can control pro-oxidant enzymes like nitric oxide synthase and antioxidant enzymes like superoxide dismutase (Darwesh et al., 2021).

According to Lordan et al., 2020, PUFAs also promote the synthesis of endothelial nitric oxide synthase (eNOS), inhibit the expression of TGF-beta, enhance the production of advantageous prostaglandins, block the activity of angiotensin-converting enzymes, decrease the production of angiotensin II, and strengthen the parasympathetic nervous system. Because of their capacity to act on the platelet membrane via COX-1 and 12-LOX to reduce platelet aggregation and TX release and metabolize fatty acids in platelets to the advantageous oxylipine group, the PUFA omega-3, EPA, and DHA groups play a major role in the regulation of platelet function in hemostasis and thrombosis.

- Fermented foods

Morinda citrifolia and *Carica papaya* are sources of fermented dietary supplements that have been shown to have anti-inflammatory, redox balancing, immunological modulation, faster regeneration, and memory-improving properties (Kharaeva et al., 2022). An RCT has demonstrated that this combination of vitamins can alleviate the clinical symptoms of Parkinson's disease (PCS), which include weakness, diminished mental and physical functioning, headaches, dizziness, insomnia, and insomniac episodes.

Treatment for psychological and mental health conditions also includes addressing apathy, anxiety, melancholy, and emotional instability (anger, irritability, sadness). Additionally, by lowering tachycardia, arrhythmia, and heart discomfort, this vitamin shields the heart. It also affects the bones, joints, and muscles, which helps prevent discomfort in those areas. In fact, it lessens hair loss and dry skin (Zhang K et al., 2022).

6. Conclusion

Studies on complementary and alternative therapies, including herbal remedies, conventional medicine, nutraceuticals and supplements, have demonstrated promising outcomes in treating problems that affect PCS patients. This is demonstrated by the patients' improved clinical conditions with no appreciable side effects. Because of this, these modalities represent a potentially effective alternative therapy for PCS patients. Although there are positive prospects for the combination of these alternative therapeutic approaches, there is still uncertainty regarding the safety and efficacy of the combined therapy due to potential side effects. In selecting alternative therapies for PCS patients, several factors should be considered, such as the patients' psychosocial, the alternative therapies' cost-effectiveness, and the clinical symptoms in the patients. It must be considered in order for medical professionals to identify and offer appropriate alternative therapies in order to treat PCS patients' symptoms as effectively as possible.

Acknowledgements

We would like to extend their gratitude first to Bambang Haryo Tri Satya Wahyono for giving support on this paper and most especially in encouraging authors to finish this paper.

References

- Al-Buhadily AK, Hussien NR, Al-Niemi MS, Al-Kuraishy HM, Al-Gareeb AI., 2021. Misfortune and spy story in the neurological manifestations of Covid-19, *J Pak Med Assoc.*, 71(12), p. S157–60.
- Albergamo A, Apprato G, Silvagno F., 2023. The Role of Vitamin D in Supporting Health in the COVID- 19 Era, *Int J Mol Sci*, 23(7).
- Alemanno F, Houdayer E, Parma A, Spina A, Del Forno A, Scatolini A, et al., 2021. COVID-19 cognitive deficits after respiratory assistance in the subacute phase: A COVID rehabilitation unit experience, *PLoS One* [Internet], 16(2 February), p. 1–12. Available from: <http://dx.doi.org/10.1371/journal.pone.0246590>.
- Anderson BJ, Jurawanichkul S, Kligler BE, Marantz PR, Evans R., 2019. Interdisciplinary Relationship Models for Complementary and Integrative Health: Perspectives of Chinese Medicine Practitioners in the United States, *J Altern Complement Med*, 2019, 25(3), p. 288–95.
- Balkrishna A, Pokhrel S, Singh J, Varshney A., 2020. Withanone from *Withania somnifera* May Inhibit Novel Coronavirus (COVID-19) Entry by Disrupting Interactions between Viral S-Protein Receptor Binding Domain and Host ACE2 Receptor, Available from: <https://doi.org/10.21203/rs.3.rs-17806/v1>.
- Barrea L, Grant WB, Frias-Toral E, Vetrani C, Verde L, de Alteriis G, et al., 2022. Dietary Recommendations for Post- COVID-19 Syndrome, *Nutrients*, 14(6), p. 1–15.
- Basu-Ray I, Metri K, Khanra D, Revankar R, Chinnaiyan KM, Raghuram N, et al., 2022. A narrative review on yoga: a potential intervention for augmenting immunomodulation and mental health in COVID-19, *BMC*

- Complement Med Ther [Internet], 22(1), p. 1–13. Available from: <https://doi.org/10.1186/s12906-022-03666-2>.
- Batiha GES, Al-kuraishy HM, Al-Gareeb AI, Welson NN., 2022. Pathophysiology of Post-COVID syndromes: a new perspective, Virol J [Internet],;19(1), p. 1–20. Available from: <https://doi.org/10.1186/s12985-022-01891-2>
- Boehmer AA, Georgopoulos S, Nagel J, Rostock T, Bauer A, Ehrlich JR., 2020. Acupuncture at the auricular branch of the vagus nerve enhances heart rate variability in humans: An exploratory study, Hear Rhythm O2 [Internet], 1(3), p. 215–21. Available from: <http://dx.doi.org/10.1016/j.hroo.2020.06.001>
- Bushell W, Castle R, Williams MA, Brouwer KC, Tanzi RE, Chopra D, et al., 2020. Meditation and Yoga Practices as Potential Adjunctive Treatment of SARS-CoV-2 Infection and COVID-19: A Brief Overview of Key Subjects, J Altern Complement Med., 26(7), p. 547–56.
- Cañas CA., 2020. The triggering of post COVID-19 autoimmunity phenomena could be associated with both transient immunosuppression and an inappropriate form of immune reconstitution in susceptible individuals, Med Hypotheses [Internet], 145(August), p. 110345. Available from: <https://doi.org/10.1016/j.mehy.2020.110345>
- Cao JF, Gong Y, Wu M, Xiong L, Chen S, Huang H, et al., 2022. Molecular docking and molecular dynamics study Lianhua Qingwen granules (LHQW) treats COVID-19 by inhibiting inflammatory response and regulating cell survival, Front Cell Infect Microbiol., 12(November), p. 1–18.
- Carvalho-schneider C, Laurent E, Lemaignan A, Laribi S, Flament T, Beau E, et al., 2020. Follow-up of adults with noncritical COVID-19 two months after symptom onset, Clin Microbiol Infect., 27(January), p. 258–63.
- Castanares-Zapatero D, Chalon P, Kohn L, Dauvrin M, Detollenaere J, Maertens de Noordhout C, et al., 2022. Pathophysiology and mechanism of long COVID: a comprehensive review, Ann Med [Internet], 54(1), p. 1473–87. Available from: <https://doi.org/10.1080/07853890.2022.2076901>.
- Catalano A, Iacopetta D, Ceramella J, Maio AC De, Basile G, Giuzio F, et al., 2022. Are Nutraceuticals Effective in COVID- 19 and Post-COVID Prevention and Treatment?, Foods, 11(18), p. 1–26.
- Chen C, Hupert SR, Zimmermann L, Shi X, Fritsche LG, Mukherjee B., 2022. Global Prevalence of Post COVID-19 Condition or Long COVID: A Meta-Analysis and Systematic Review 2 Running Title: Post COVID-19 Condition Meta- Analysis 3 4, J Infect Dis [Internet], (jiac136), p. 1–32. Available from: https://academic.oup.com/journals/pages/open_access/funder_policies/chorus/standard_publication_mode
- Chien TJ, Liu CY, Chang YI, Fang CJ, Pai JH, Wu YX, et al., 2022, Therapeutic effects of herbal- medicine combined therapy for COVID-19: A systematic review and meta-analysis of randomized controlled trials, Front Pharmacol., 13(September), p. 1–17.
- Cruz J, Trombley J, Carrington L, Cheng X., 2021. Properties of the Novel Chinese Herbal Medicine Formula Qu du Qiang Fei i Hao Fang Warrant Further Research to Determine Its Clinical Efficacy in COVID-19 Treatment, Med Acupunct., 33(1), p. 71–82.

- Darwesh AM, Bassiouni W, Sosnowski DK, Seubert JM., 2021. Can N-3 polyunsaturated fatty acids be considered a potential adjuvant therapy for COVID-19-associated cardiovascular complications?, *Pharmacol Ther J.*, 219(January), p. 107703.
- Dhane RM, Shinde RB., 2022. Herbal Remedies in Post-COVID Trauma : A Systematic Review, 19(December), p. 875–9.
- Driscoll L., 2020. Extracellular vesicles from mesenchymal stem cells as a Covid-19 treatment, *Drug Discov Today*, 25(1), p. 1124–5.
- Easwaran K, Gopalasingam Y, Green DD, Lach V, Melnyk JA, Wan C, et al., 2021. Effectiveness of Tai Chi for health promotion for adults with health conditions: a scoping review of Meta-analyses, *Disabil Rehabil* [Internet], 43(21), p. 2978– 89. Available from: <https://doi.org/10.1080/09638288.2020.1725916>.
- Fratta Pasini AM, Stranieri C, Cominacini L, Mozzini C., 2021. Potential role of antioxidant and anti-inflammatory therapies to prevent severe sars-cov-2 complications, *Antioxidants*, 10(2), p. 1–22.
- Fernandes AL, Murai IH, Reis BZ, Sales LP, D M, Santos, et al., 2018. Effect of a Single High Dose of Vitamin D3 on Cytokines, Chemokines, and Growth Factor in Patients With Moderate to Severe COVID- 19, *Am J Epidemiol.*, 187(4)(June), p. 717-25.
- Fuchs B, Sjöberg L, Westerberg CM, Ekoff M, Swedin L, Dahlén SE, et al., 2012. Mast cell engraftment of the peripheral lung enhances airway hyperresponsiveness in a mouse asthma model, *Am J Physiol - Lung Cell Mol Physiol*, 303(12), p. 1027–36.
- Goërtz YMJ, Van Herck M, Delbressine JM, Vaes AW, Meys R, Machado FVC, et al., 2020. Persistent symptoms 3 months after a SARS-CoV-2 infection: the post COVID-19 syndrome?, *ERJ Open Res* [Internet], 6(4), p. 00542–2020. Available from: <http://dx.doi.org/10.1183/23120541.00542-2020>
- Hampton A, Bartz M., 2021. Therapeutic Efficacy of Yoga for Common Primary Care Conditions, *Wis Med J.*, 120(4), p. 293–300.
- Helianthi DR, Barasila AC, Harris S, Sinto R, Khaedir Y, Irman., 2022. The Adjunctive Effects of Acupuncture for Hospitalized COVID-19 Patients: a Single-Blinded Randomized Clinical Study, *JAMS J Acupunct Meridian Stud.*, 15(4), p. 247–54.
- Hope AA, Evering TH., 2022. Postacute Sequelae of Severe Acute Respiratory Syndrome Coronavirus 2 Infection, *Infect Dis Clin North Am* [Internet], 6(2), p. 379–95, Available from: <https://doi.org/10.1016/j.idc.2022.02.004>
- Jamshidi E, Babajani A, Soltani P, Niknejad H., 2021. Proposed Mechanisms of Targeting COVID-19 by Delivering Mesenchymal Stem Cells and Their Exosomes to Damaged Organs, *Stem Cell Rev Reports.*, 17(1), p. 176 –92.
- Jia H, Han Z, Zhang K, Tang Q, Sun K, Huang H, et al., 2020. Acupuncture and related interventions for anxiety in coronavirus disease 2019: A protocol for systematic review and meta-analysis, *Medicine (Baltimore)*, 2020, 99(30), p. 21317.

- Jones R, Davis A, Stanley B, Julious S, Ryan D, Jackson DJ, et al., 2021. Risk Predictors and Symptom Features of Long COVID Within a Broad Primary Care Patient Population Including Both Tested and Untested Patients, *Pragmatic Obs Res.*, 12(July), p. 93–104.
- Kharaeva Z, Shokarova A, Shomakhova Z, Ibragimova G, Trakhtman P, Trakhtman I, et al., 2022. Fermented Carica papaya and Morinda citrifolia as Perspective Food Supplements for the Treatment of Post-COVID Symptoms: Randomized Placebo- Controlled Clinical Laboratory Study, *Nutrients*, 14(11).
- Kim TH, Jeon SR, Kang JW, Kwon S., 2022. Complementary and Alternative Medicine for Long COVID: Scoping Review and Bibliometric Analysis, *Evidence-based Complement Altern Med*, 2022.
- Koc HC, Xiao J, Liu W, Li Y, Chen G., 2022. Long COVID and its Management, *Int J Biol Sci*, 18(12), p. 4768–80.
- Koné-Paut I, Cimaz R., 2020. Is it Kawasaki shock syndrome, Kawasaki-like disease or pediatric inflammatory multisystem disease? The importance of semantic in the era of COVID-19 pandemic, *RMD Open*, 6(2), p. 1–4.
- Kumar S, Malaiya A, Mishra G, Jain D, Kesharwani P., 2020. An exhaustive comprehension of the role of herbal medicines in Pre- and Post-COVID manifestations, *J Ethnopharmacol*, 296(January), p. 115420.
- Kwon S, Jung WS, Yang S, Jin C, Cho SY, Park SU, et al., 2018. Safety of acupuncture in patients taking newer oral anticoagulants: A retrospective chart review study, *Evidence-based Complement Altern Med*, 2018, p. 1–5.
- Li J, Zhang K, Bao J, Yang J, Wu C., 2022. Potential mechanism of action of Jing Fang Bai Du San in the treatment of COVID-19 using docking and network pharmacology, *Int J Med Sci.*, 19(2), p. 213–24.
- Li L, Zhang Z, Zhou W, Chen J, Jin H., 2020. Lianhua Qingwen prescription for Coronavirus disease 2019 (COVID-19) treatment: Advances and prospects, *Biomed Pharmacother*, 130(January), p. 110641.
- Liu K, Zhang W, Yang Y, Zhang J, Li Y, Chen Y., 2020. Respiratory rehabilitation in elderly patients with COVID-19: A randomized controlled study, *Complement Ther Clin Pract.*, 39(January), p. 101166.
- Liu YS, Liu Q, Jiang YL, Yang WT, Huang H Bin, Shi CW, et al., 2020. Surface-Displayed Porcine IFN- λ 3 in Lactobacillus plantarum Inhibits Porcine Enteric Coronavirus Infection of Porcine Intestinal Epithelial Cells, *J Microbiol Biotechnol.*, 30(4), p. 515–25.
- Lordan R, Redfern S, Tsoupras A, Zabetakis I., 2020. Inflammation and cardiovascular disease: Are marine phospholipids the answer?, *Food Funct.*, 11(4), p. 2861–85.
- Macedo S, Ribeiro F, Ponte S, Soares P, Boaventura P., 2023. Yoga for COVID-19: An ancient practice for a new condition – A literature review, *Complement Ther Clin Pract.*, 50(January), p. 101717.
- Magdalena A, Pop A, Iordache F, Stanca L., 2022. Antioxidant, anti-inflammatory and immunomodulatory roles of vitamins in COVID-19 therapy, *Eur J Med Chem.*, 232(January), p. 114175.
- Maurya DK, Sharma D., 2022. Evaluation of traditional ayurvedic Kadha for prevention and management of the novel Coronavirus (SARS-CoV-2) using in silico approach, *J Biomol Struct Dyn.*, 40(9), p. 3949–64.

- Masnadi Shirazi K, Nikniaz Z, Masnadi Shirazi A, Rohani M., 2018. Vitamin A supplementation decreases disease activity index in patients with ulcerative colitis: A randomized controlled clinical trial, *Complement Ther Med* [Internet], 41(September), p. 215–9. Available from: <https://doi.org/10.1016/j.ctim.2018.09.026>
- Maxwell AJ, Ding J, You Y, Dong Z, Chehade H, Alvero A, et al., 2021. Identification of key signaling pathways induced by SARS-CoV2 that underlie thrombosis and vascular injury in COVID-19 patients, *J Leukoc Biol.*, 109(1), p. 35–47.
- Mehandru S, Merad M., 2022. Pathological sequelae of long-haul COVID, *Nat Immunol.*, 23(2), p. 194– 202.
- Mitrani RD, Dabas N, Goldberger JJ., 2020. COVID-19 cardiac injury: Implications for long-term surveillance and outcomes in survivors, *Hear Rhythm.*, 17(11), p. 1985–90.
- Monari P, Gualdi G, Bettoni G, Costa R, Ragni G, Zani F, et al., 2022. Post-SARS-CoV-2 Acute Telogen Effluvium: An Expected Complication, *J Clin Med*, 11(5).
- Moreno-Perez O, Merino E, Alfayate R, Torregrosa ME, Andres M, Leon-Ramirez JM, et al. 2022. Male pituitary–gonadal axis dysfunction in post-acute COVID-19 syndrome—Prevalence and associated factors: A Mediterranean case series. *Clin Endocrinol (Oxf)*, 96(3):353–62.
- Morita A, Murakami A, Uchihara T, Ohashi N, Ryu K, Watanabe Y, et al., 2022. Case Report: Acupuncture is an effective treatment for olfactory dysfunction in the post COVID-19 condition, *Front Neurol.*, 13.
- Mukherjee R, Bhattacharya A, Bojkova D, Mehdi pour AR, Shin D, Khan KS, et al., 2021. Famotidine inhibits toll-like receptor 3-mediated inflammatory signaling in SARS-CoV-2 infection, *J Biol Chem* [Internet], 297(2), p. 100925. Available from: <https://doi.org/10.1016/j.jbc.2021.100925>
- Naureen Z, Dautaj A, Nodari S, Fioretti F, Dhuli K, Anpilogov K, et al., 2021. Proposal of a food supplement for the management of post-COVID syndrome, *Eur Rev Med Pharmacol Sci*, 25(1), p. 67–73.
- O’Mahoney LL, Routen A, Gillies C, Ekezie W, Welford A, Zhang A, et al., 2023. The prevalence and long-term health effects of Long Covid among hospitalised and non-hospitalised populations: A systematic review and meta-analysis, *eClinicalMedicine*, 55, p. 1–10.
- Osmanov IM, Spiridonova E, Bobkova P, Gamirova A, Shikhaleva A, Andreeva M, et al., 2022. Risk factors for long covid in previously hospitalised children using the ISARIC Global follow-up protocol: A prospective cohort study, *Eur Respir J* [Internet], 59(2). Available from: <http://dx.doi.org/10.1183/13993003.01341-2021>
- Pal A, Squitti R, Picozza M, Pawar A, Rongioletti M, Dutta AK, et al., 2021. Zinc and COVID-19: Basis of Current Clinical Trials, *Biol Trace Elem Res*, 199(8), p. 2882–92.
- Panda AK, Kar S, Rai AK, Rao BCS, Srikanth N., 2021. AYUSH- 64: A potential therapeutic agent in COVID-19, *J Ayurveda Integr Med* [Internet], 13(2), p. 100538. Available from: <https://doi.org/10.1016/j.jaim.2021.100538>.
- Pastor N, Collado MC, Manzoni P., 2021. Phytonutrient and nutraceutical action against COVID-19: Current review of characteristics and benefits, *Nutrients*, 13(2), p. 1–10.

- Polero P, Rebollo-Seco C, Adsuar JC, Pérez-Gómez J, Rojo-Ramos J, Manzano-Redondo F, et al., 2021. Physical activity recommendations during COVID-19: Narrative review, *Int J Environ Res Public Health*, 18(1), p. 1–24.
- Popescu MN, Berteau M, Beiu C, Popa LG, Mihai MM, Iliescu MG, et al., 2022. Complementary Strategies to Promote Hair Regrowth in Post COVID-19 Telogen Effluvium, *Clin Cosmet Investig Dermatol*, 15(March), p. 735–43.
- Range K, M D, Moser YA., 2013. Sepsis-induced immunosuppression: from cellular dysfunctions to immunotherapy, *Nat Rev Immunol.*, 23(12), p. 862–74.
- Rastogi S, Pandey DN, Singh RH., 2022. COVID-19 pandemic: A pragmatic plan for ayurveda intervention, *J Ayurveda Integr Med* [Internet], 13(1), p. 100312. Available from: <https://doi.org/10.1016/j.jaim.2020.04.002>.
- Rohani M, Mozaffar H, Mesri M, Shokri M, Delaney D, Karimy M., 2022. Evaluation and comparison of vitamin A supplementation with standard therapies in the treatment of patients with COVID-19, *East Mediterr Heal J*, 28(9), p. 673–81.
- Rossato MS, Brilli E, Ferri N, Giordano G, Tarantino G., 2021. Observational study on the benefit of a nutritional supplement, supporting immune function and energy metabolism, on chronic fatigue associated with the SARS-CoV-2 post-infection progress, *Clin Nutr ESPEN*, 46(January), p. 510–8.
- Runfeng L, Yunlong H, Jicheng H, Weiqi P, Qin Hai M, Yongxia S., 2020. Lianhuaqingwen exerts anti- viral and anti- inflammatory activity against novel coronavirus (SARS-CoV-2), *Pharmacol Res J*, 156(January), p. 104761.
- Saeed SA, Cunningham K, Bloch RM., 2019. Depression and anxiety disorders: Benefits of exercise, yoga, and meditation, *Am Fam Physician*, 99(10), p. 620–7.
- Schweiger V, Secchettin E, Castellani C, Martini A, Mazzocchi E, Picelli A, et al., 2020. Comparison between Acupuncture and Nutraceutical Treatment with Migratens® in Patients with Fibromyalgia Syndrome: A Prospective Randomized Clinical Trial, *Nutrients*, 12(3).
- Scoppettuolo P, Borrelli S, Naeije G., 2020. Neurological involvement in SARS-CoV-2 infection: A clinical systematic review, *Brain, Behav Immun - Heal* [Internet], 5(June), p. 100094. Available from: <https://doi.org/10.1016/j.bbih.2020.100094>
- Shree P, Mishra P, Selvaraj C, Singh SK, Chaube R, Garg N, et al., 2022. Targeting COVID-19 (SARS- CoV-2) main protease through active phytochemicals of ayurvedic medicinal plants—Withania somnifera (Ashwagandha), Tinospora cordifolia (Giloy) and Ocimum sanctum (Tulsi)—a molecular docking study, *J Biomol Struct Dyn* [Internet], 40(1), p. 190–203. Available from: <https://doi.org/10.1080/07391102.2020.1810778>
- Shu C, Feng S, Cui Q, Cheng S, Wang Y., 2021. Impact of Tai Chi on CRP, tnf-alpha and il-6 in inflammation: A systematic review and meta-analysis, *Ann Palliat Med.*, 10(7), p. 7468–78.
- Singh R, Shaik L, Mehra I, Kashyap R, Surani S., 2021. Novel and Controversial Therapies in COVID- 19, *Open Respir Med J*, 14(1), p. 79–86.

- Song G, Fiocchi C, Achkar JP., 2019. Acupuncture in inflammatory bowel disease, *Inflamm Bowel Dis.*, 25(7), p. 1129–39.
- Srinivasan, V.; Kandakurti, P. K.; Alagesan, J.; Suganthirababu, P.; Kishore Jebasingh, T.; Jenifer Augustina, S.; Anitha, A., 2021. Efficacy of pursed lip breathing with bhastrika pranayama vs incentive spirometry in rehabilitating post Covid 19 follow up-a randomized control study, *Turkish Journal of Physiotherapy and Rehabilitation*, 32(3), p. 402-407.
- Sugimoto MA, Sousa LP, Pinho V, Perretti M, Teixeira MM., 2016. Resolution of inflammation: What controls its onset?, *Front Immunol.*, 7(APR).
- Sugiyama A, Miwata K, Kitahara Y, Okimoto M, Abe K, E B, et al., 2022. Long COVID occurrence in COVID-19 survivors, *Sci Rep [Internet]*, 12(1), p. 1–11. Available from: <https://doi.org/10.1038/s41598-022-10051-z>.
- Tosato M, Ciciarello F, Zazzara MB, Pais C, Saveria G, Picca A, et al., 2022. Nutraceuticals and Dietary Supplements for Older Adults with Long COVID-19, *Clin Geriatr Med.*, 38(January), p. 565–91.
- Trager RJ, Brewka EC, Kaiser CM, Patterson AJ, Dusek JA., 2022. Acupuncture in Multidisciplinary Treatment for Post- COVID-19 Syndrome, *Med Acupunct.*, 34(3), p. 177–83.
- Visco V, Vitale C, Rispoli A, Izzo C, Virtuoso N, Ferruzzi GJ, et al., 2022. Post COVID-19 Syndrome: Involvement and Interactions between Respiratory, Cardiovascular and Nervous Systems, *J Clin Med.*, 11(3).
- Wang, R. Q., Liu, J. X., Zhang, Z. D., Wen, J., Han, P., Wu, H. H., Jia, Y. J., Jia, C. S., & Pan, L. J., 2020. Zhen ci yan jiu = Acupuncture research, 45(5), p. 345–350. <https://doi.org/10.13702/j.1000-0607.200275>
- Wang S, Su K, Pariente 2021. CM., The three frontlines against COVID-19: Brain, Behavior, and Immunity, *Brain Behav Immun.*, 93(January), p. 409–14.
- Wang Y, Greenhalgh T, Wardle J., 2022. Chinese herbal medicine (“3 medicines and 3 formulations”) for COVID-19: rapid systematic review and meta-analysis, *J Eval Clin Pract*, 28(1), p. 13–32.
- Wang Y, Jin X, Fan Q, Li C, Zhang M, Wang Y, et al., 2022. Deciphering the Active Compounds and Mechanisms of HSBDF for Treating ALI via Integrating Chemical Bioinformatics Analysis, *Front Pharmacol.*, 13(June), p. 1–15.
- Wu K, Li Y, Zou Y, Ren Y, Wang Y, Hu X, et al., 2022. Tai Chi increases functional connectivity and decreases chronic fatigue syndrome: A pilot intervention study with machine learning and fMRI analysis, *PLoS One [Internet]*, 17(12), p. e0278415. Available from: <http://dx.doi.org/10.1371/journal.pone.0278415>.
- Xia QD, Xun Y, Lu JL, Lu YC, Yang YY, Zhou P, et al., 2020. Network pharmacology and molecular docking analyses on Lianhua Qingwen capsule indicate Akt1 is a potential target to treat and prevent COVID-19, *Cell Prolif.*, 53(12), p. 1–13.
- Xiong L, Cao J, Yang X, Chen S, Wu M, Wang C, et al., 2022. Exploring the mechanism of action of Xuanfei Baidu granule (XFBD) in the treatment of COVID-19 based on molecular docking and molecular dynamics, *Front Cell Infect Microbiol.*, 12(August), p. 1–21.

- Xiong Y, Li NX, Duan N, Liu B, Zhu H, Zhang C, et al., 2020. Traditional Chinese Medicine in Treating Influenza: From Basic Science to Clinical Applications, *Front Pharmacol*, 11(September), p. 1–9.
- Xu Y, Yang L, Wang L, Chen F., 2021. Potential therapeutic effect of Shufeng Jiedu capsule and its major herbs on coronavirus disease 2019 (COVID-19): A review, *Drug Discov Ther.*, 15(6), p. 289–99.
- Xu YF, Zhuang WY, Mao M, et al., 2021. Discussion on the application of acupuncture and moxibustion therapy in the prevention and treatment of ancient epidemic disease (in Chinese), *China J Tradit Chin Med Pharm.*, 36(05), p. 2464–2467.
- Yates CR, Bruno EJ, Yates MED., 2022. *Tinospora Cordifolia*: A review of its immunomodulatory properties, *J Diet Suppl.*, 19(2), p. 271–85. Available from: <https://doi.org/10.1080/19390211.2021.1873214>.
- Zhang J, Rao X, Li Y, Zhu Y, Liu F, Guo G, et al., 2021. Pilot trial of high-dose vitamin C in critically ill COVID-19 patients, *Ann Intensive Care* [Internet], 11(1), p. 3–14. Available from: <https://doi.org/10.1186/s13613-020-00792-3>.
- Zhang J, Zhang Z, Huang S, Qiu X, Lao L, Huang Y, et al., 2022. Acupuncture for cancer-related insomnia: A systematic review and meta-analysis, *Phytomedicine* [Internet], 102(December 2021), p. 154160. Available from: <https://doi.org/10.1016/j.phymed.2022.154160>.
- Zhang Q, Xu X, Sun S, Cao F, Li J, Qi X, et al., 2020. Efficacy of acupuncture and moxibustion in adjuvant treatment of patients with novel coronavirus disease 2019 (COVID-19): A protocol for systematic review and meta analysis. *Medicine (Baltimore)*, 99(28), p. 21039.
- Zhang S, Zou L, Chen LZ, Yao Y, Loprinzi PD, Siu PM, et al., 2019. Effect of tai chi chuan on negative emotions in non- clinical populations: A meta-analysis and systematic review, *Int J Environ Res Public Health*, 16(17).
- Zhao B, Ling Y, Li J, Peng Y, Huang J, Wang Y, et al., 2021. Beneficial aspects of high dose intravenous vitamin C on patients with COVID-19 pneumonia in severe condition: a retrospective case series study, *Ann Palliat Med*, 10(2), p. 1599– 609.
- Zhao C, Li L, Yang W, Lv W, Wang J, Guo J, et al., 2021. Chinese Medicine Formula Huashibaidu Granule Early Treatment for Mild COVID-19 Patients: An Unblinded, Cluster-Randomized Clinical Trial, *Front Med.*, 8(September), p. 1–10.
- Zhao J, Tian S, Lu D, Yang J, Zeng H, Zhang F, et al., 2021. Systems pharmacological study illustrates the immune regulation, anti-infection, anti-inflammation, and multi-organ protection mechanism of Qing-Fei-Pai-Du decoction in the treatment of COVID-19, *Phytomedicine* [Internet], 85(August 2020), p. 153315. Available from: <https://doi.org/10.1016/j.phymed.2020.153315>
- Zhang K, Meng J, Li X, Tang X, Ma S, Lv Y, et al., 2022. Noni (*Morinda citrifolia* L.) wine prevents the oxidative stress and obesity in mice induced by high fat diet, *J Food Biochem.*, 44(11), p. 1–14.
- Zhu Y, Luo L, Zhang M, Song X, Wang P, Zhang H., 2023. Xuanfei Baidu Formula attenuates LPS- induced acute lung injury by inhibiting the NF- κ B signaling pathway, *J Ethnopharmacol.*, 301(January), p. 115833.

Zou L, Yeung A, Quan X, Hui SSC, Hu X, Chan JSM, et al., 2018. Mindfulness-based baduanjin exercise for depression and anxiety in people with physical or mental illnesses: A systematic review and meta-analysis, *Int J Environ Res Public Health*, 15(2).